

Counting topologies of metric holomorphic polynomial field with simple zeros


Art Duval¹, Martín Eduardo Frías-Armenta²

¹University of Texas at El Paso, ²Universidad de Sonora


AMS Southeastern Sectional Meeting
Special Session on Geometric and Topological Combinatorics
University of Florida
November 2, 2019

AD supported by Simons Foundation Grant 516801

Setting the scene: Trees from flow diagrams




Metric holomorphic polynomial field with simple zeros




Phase portrait of $X(z) = 2i - iz - 2iz^4 + iz^5 \frac{\partial}{\partial z}$

Metric holomorphic polynomial field with simple zeros




Phase portrait of $X(z) = 2i - iz - 2iz^4 + iz^5 \frac{\partial}{\partial z}$

Complex rotation




Phase portrait of $X(z)(\frac{1}{2} - i)$

Complex rotation



Phase portrait of $X(z)(-1 + \frac{i}{2})$

Put it all together, and get a graph



So we are looking at unlabeled trees with black and white vertices

- ▶ no white vertices are adjacent to each other
- ▶ each white vertex is adjacent to at least three black vertices
- ▶ no restriction on neighbors of black vertices

Trees




So we are looking at unlabeled trees with black and white vertices

- ▶ no white vertices are adjacent to each other
- ▶ each white vertex is adjacent to at least three black vertices
- ▶ no restriction on neighbors of black vertices


We want to count such trees up to rotation (but not reflection)

Example

The first two are the same, but the third is different.






Flashback: Counting (unlabeled) trees



How to grow different kinds of rooted trees, recursively

- ▶ Rooted trees:
 - ▶ $\mathcal{A} = X \cdot E(\mathcal{A})$,
 - ▶ E stands for “set of”
- ▶ Ordered rooted tree:
 - ▶ $\mathcal{A}_L = X \cdot L(\mathcal{A}_L)$
 - ▶ L stands for “linear order”
- ▶ Planar rooted trees:
 - ▶ $P = X + X \cdot C(\mathcal{A}_L)$
 - ▶ C stands for “cyclic order”

Example



Unrooting I: Center of tree



Definition

Center of a tree is the set of vertices v that minimize

$$\max_u d(u, v)$$

It is always either a single vertex, or an edge.

Example



Unrooting I: Center of tree



Definition

Center of a tree is the set of vertices v that minimize

$$\max_u d(u, v)$$

It is always either a single vertex, or an edge. So this **naturally roots a tree** at either a vertex or an edge.

Example



Unrooting II: Dissymmetry theorem

Theorem (Dissymmetry)

$$\mathcal{A} + E_2(\mathcal{A}) = \mathfrak{a} + \mathcal{A}^2,$$

where \mathfrak{a} denotes unrooted trees and E_2 is the species of sets with exactly two elements.

Unrooting II: Dissymmetry theorem

Theorem (Dissymmetry)

$$\mathcal{A} + E_2(\mathcal{A}) = \mathfrak{a} + \mathcal{A}^2,$$

where \mathfrak{a} denotes unrooted trees and E_2 is the species of sets with exactly two elements.

Proof.

(Sketch) LHS is trees rooted at a vertex or an edge. RHS is trees (unrooted) or ordered pair of rooted trees.

Unrooting II: Dissymmetry theorem


Theorem (Dissymmetry)

$$\mathcal{A} + E_2(\mathcal{A}) = \mathfrak{a} + \mathcal{A}^2,$$

where \mathfrak{a} denotes unrooted trees and E_2 is the species of sets with exactly two elements.

Proof.

(Sketch) LHS is trees rooted at a vertex or an edge. RHS is trees (unrooted) or ordered pair of rooted trees. So we need isomorphism between trees rooted at vertex or edge **other** than the center, with ordered pairs of rooted trees. \square



Quick note about unlabeled


Theorem (Dissymmetry)

$$\mathcal{A} + E_2(\mathcal{A}) = \mathfrak{a} + \mathcal{A}^2,$$

where \mathfrak{a} denotes unrooted trees and E_2 is the species of sets with exactly two elements.


Dissymmetry theorem allows us to count unrooted, but still labeled trees. To unlabeled the trees, we need “cycle index series”.

Return to the present day: Counting our trees



Black and white vertices, not at the root

Similar to ordered rooted trees, but now color-aware



$$Y_1 = X_1 \cdot L(Y_1 + Y_2) \quad Y_2 = X_2 \cdot L_{\geq 2}(X_1 \cdot L(Y_1 + Y_2))$$

$$Y_3 = Y_1 + Y_2 = X_1 \cdot L(Y_3) + X_2 \cdot L_{\geq 2}(X_1 \cdot L(Y_3))$$

Recursive equation

$$Y_3 = Y_1 + Y_2 = X_1 \cdot L(Y_3) + X_2 \cdot L_{\geq 2}(X_1 \cdot L(Y_3))$$

$$y_3 = x_1 \ell + x_2 \frac{(x_1 \ell)^2}{1 - (x_1 \ell)}$$

where $\ell = \frac{1}{1-y_3}$.

Recursive equation

$$Y_3 = Y_1 + Y_2 = X_1 \cdot L(Y_3) + X_2 \cdot L_{\geq 2}(X_1 \cdot L(Y_3))$$

$$y_3 = x_1 \ell + x_2 \frac{(x_1 \ell)^2}{1 - (x_1 \ell)}$$

where $\ell = \frac{1}{1-y_3}$. Simplifying,

$$x_1 + x_1^2(x_2 - 1) - (y_3 - 1)^2 y_3 - x_1 y_3^2 = 0.$$

Unique real root $y_3(x_1, x_2) =$

Recursive equation

$$Y_3 = Y_1 + Y_2 = X_1 \cdot L(Y_3) + X_2 \cdot L_{\geq 2}(X_1 \cdot L(Y_3))$$

$$y_3 = x_1 \ell + x_2 \frac{(x_1 \ell)^2}{1 - (x_1 \ell)}$$

where $\ell = \frac{1}{1-y_3}$. Simplifying,

$$x_1 + x_1^2(x_2 - 1) - (y_3 - 1)^2 y_3 - x_1 y_3^2 = 0.$$

Unique real root $y_3(x_1, x_2) =$

$$\frac{\frac{2-x_1}{3} + (2^{1/3}(-1+4x_1-x_1^2))}{\left(3\left(2-12x_1+15x_1^2+2x_1^3-27x_1^2x_2+\sqrt{4(-1+4x_1-x_1^2)^3+(2-12x_1+15x_1^2+2x_1^3-27x_1^2x_2)^2}\right)\right)^{1/3}}$$
$$-\frac{1}{3 \cdot 2^{1/3}} \left(2-12x_1+15x_1^2+2x_1^3-27x_1^2x_2+\sqrt{4(-1+4x_1-x_1^2)^3+(2-12x_1+15x_1^2+2x_1^3-27x_1^2x_2)^2}\right)$$

Dissymmetry again

Recall

$$\mathcal{A} + E_2(\mathcal{A}) = \mathfrak{a} + \mathcal{A}^2,$$

The same arguments apply. But now, paying attention to color,

$$\mathcal{A}_R = (X_1 \cdot (1 + C(Y_3))) + (X_2 \cdot C_{\geq 3}(X_1 \cdot L(Y_3)))$$

Dissymmetry again

Recall

$$\mathcal{A} + E_2(\mathcal{A}) = \mathfrak{a} + \mathcal{A}^2,$$

The same arguments apply. But now, paying attention to color,

$$\begin{aligned}\mathcal{A}_R &= (X_1 \cdot (1 + C(Y_3))) + (X_2 \cdot C_{\geq 3}(X_1 \cdot L(Y_3))) \\ E_2(\mathcal{A}_R) &= E_2(Y_1) + Y_2 \cdot Y_1 = E_2(Y_3) - E_2(Y_2)\end{aligned}$$

Dissymmetry again

Recall

$$\mathcal{A} + E_2(\mathcal{A}) = \mathfrak{a} + \mathcal{A}^2,$$

The same arguments apply. But now, paying attention to color,

$$\mathcal{A}_R = (X_1 \cdot (1 + C(Y_3))) + (X_2 \cdot C_{\geq 3}(X_1 \cdot L(Y_3)))$$

$$E_2(\mathcal{A}_R) = E_2(Y_1) + Y_2 \cdot Y_1 = E_2(Y_3) - E_2(Y_2)$$

$$\mathcal{A}_R^2 = Y_1^2 + 2Y_1Y_2 = Y_3^2 - Y_2^2$$

Dissymmetry again

Recall



$$\mathcal{A} + E_2(\mathcal{A}) = \mathbf{a} + \mathcal{A}^2,$$

The same arguments apply. But now, paying attention to color,

$$\begin{aligned}\mathcal{A}_R &= (X_1 \cdot (1 + C(Y_3))) + (X_2 \cdot C_{\geq 3}(X_1 \cdot L(Y_3))) \\ E_2(\mathcal{A}_R) &= E_2(Y_1) + Y_2 \cdot Y_1 = E_2(Y_3) - E_2(Y_2) \\ \mathcal{A}_R^2 &= Y_1^2 + 2Y_1Y_2 = Y_3^2 - Y_2^2\end{aligned}$$

This can be stated more generally for “multi-sort” species.
(And then, to remove labels, again bring in cycle index series.)

Aftermath: Data and Specializations



Data

	1	2	3	4	5	6	7	8	9	10	11	12
0	1	1	1	2	3	6	14	34	95	280	854	2694
1	0	0	1	2	5	16	48	164	559	1952	6872	24520
2	0	0	0	0	1	5	30	146	693	3108	13608	58200
3	0	0	0	0	0	0	2	20	175	1254	7752	44112
4	0	0	0	0	0	0	0	0	7	95	1125	10108
5	0	0	0	0	0	0	0	0	0	0	19	480
6	0	0	0	0	0	0	0	0	0	0	0	0

Data

	1	2	3	4	5	6	7	8	9	10	11	12
0	1	1	1	2	3	6	14	34	95	280	854	2694
1	0	0	1	2	5	16	48	164	559	1952	6872	24520
2	0	0	0	0	1	5	30	146	693	3108	13608	58200
3	0	0	0	0	0	0	2	20	175	1254	7752	44112
4	0	0	0	0	0	0	0	0	7	95	1125	10108
5	0	0	0	0	0	0	0	0	0	0	19	480
6	0	0	0	0	0	0	0	0	0	0	0	0

No white vertices:

Unlabeled plane trees.

Data


	1	2	3	4	5	6	7	8	9	10	11	12
0	1	1	1	2	3	6	14	34	95	280	854	2694
1	0	0	1	2	5	16	48	164	559	1952	6872	24520
2	0	0	0	0	1	5	30	146	693	3108	13608	58200
3	0	0	0	0	0	0	2	20	175	1254	7752	44112
4	0	0	0	0	0	0	0	0	7	95	1125	10108
5	0	0	0	0	0	0	0	0	0	0	19	480
6	0	0	0	0	0	0	0	0	0	0	0	0

No white vertices:

Unlabeled plane trees.

Minimal black vertices:

Unlabeled 3-gonal cacti with n triangles.
(Bóna, Bousquet, Labelle, Leroux, 2000)





One white vertex

	1	2	3	4	5	6	7	8	9	10	11	12
1	0	0	1	2	5	16	48	164	559	1952	6872	24520

Triangulations of an n -gon with exactly one internal vertex.

(Brown, 1964)





One white vertex

	1	2	3	4	5	6	7	8	9	10	11	12
1	0	0	1	2	5	16	48	164	559	1952	6872	24520

Triangulations of an n -gon with exactly one internal vertex.

(Brown, 1964)



Both are circular orders of (at least three) Catalan-things (ordered rooted trees or rooted triangulations).