

DISTANCE BETWEEN RANDOM POINTS IN A CUBE

A.M. Mathai, P. Moschopoulos, G. Pederzoli

1. INTRODUCTION

The pioneering work of Ghosh (1943a,b, 1949, 1951) seems to be the first published papers containing explicit expressions for the density functions of the distance between two random points in a square or in a rectangle. In these remarkable contributions the densities are derived for three distinct configurations: a rectangle, two adjacent rectangles and squares having a common diagonal. Ghosh was tackling some problems arising from sample survey studies, a study undertaken at the Indian Statistical Institute under the direction of Professor P.C. Mahalanobis. The author was mainly interested in topographic variation in statistical fields.

Sample survey is not the only field where one requires the statistical properties of a straight path across a preassigned geometrical figure. Several distance methods have been considered by Persson (1964, 1965) in two studies related to forestry. In physics, some problems of interest are to assess the length of a gamma-ray to the wall of a nuclear reactor from a given source and the distance of a sound-ray in a chamber from one reflection to the next, or the measure of a direct trajectory across a square or a cube. Horowitz (1965) considered such problems in a variety of basic geometric shapes when the radiation sources are assumed to be isotropic. Some results in this direction were obtained much earlier by Jäger (1911). Another interesting application is concerned with the traffic flow in the central area of a town with a rectangular grid as the road system, as reported in Smeed and Jeffcoate (1963). The two survey articles by Moran (1966, 1969) also contain a list of applications, and so does the book by Eilon, Watson-Gandy and Christofides (1971). Random paths across elementary geometrical shapes are also considered by Kendall and Moran (1963), Coleman (1969) and Kingman (1965, 1969).

In the present article we consider the distance between two points which are independently and uniformly distributed inside a cube of side a . A special case of this distance also describes the distance between two independent random points in a square, a case considered by Ghosh, as mentioned earlier. Furthermore, explicit forms of the exact density and arbitrary moments of the distance are

evaluated in terms of Gauss' hypergeometric functions and Appell's function F_1 which is also Lauricella function F_D for two variables. Simpler forms are shown to be available for positive integer moments of the squared distance. Special cases of the results also include the corresponding results for the distance between two random points inside a square.

2. THE DISTANCE BETWEEN TWO RANDOM POINTS INSIDE A CUBE

Consider a cube of a side a and let $P = (x_1, y_1, z_1)$ and $Q = (x_2, y_2, z_2)$ be two random points inside this cube. Randomness means that $x_j, y_j, z_j, j = 1, 2$ are mutually independently and uniformly distributed over $[0, a]$. The distance between the points P and Q is then

$$x = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}.$$

One method of deriving the density of x is to look at the density of the sum $u = x^2 = u_1 + u_2 + u_3$ where

$$u_1 = (x_2 - x_1)^2, \quad u_2 = (y_2 - y_1)^2, \quad u_3 = (z_2 - z_1)^2.$$

Then the densities of u_1, u_2, u_3 are available from elementary computations. They are

$$f_j(u_j) = \frac{1}{a^2} \left[au_j^{-\frac{1}{2}} - 1 \right], \quad 0 \leq u_j \leq a^2, \quad j = 1, 2, 3.$$

Making the transformation $u_1 = t, u_2 = v_1 - t$ the joint density of v_1 and t is

$$f(v_1, t) = f_1(t)f_2(v_1 - t) = \frac{1}{a^4} \left\{ a^2 t^{-\frac{1}{2}} (v_1 - t)^{-\frac{1}{2}} - at^{-\frac{1}{2}} - a(v_1 - t)^{-\frac{1}{2}} + 1 \right\}$$

and the marginal density of v_1 , denoted by $g_1(v_1)$, is available as

$$g_1(v_1) = \begin{cases} h_1(v_1), & 0 \leq v_1 \leq a^2 \\ b_2(v_2), & a^2 \leq v_1 \leq 2a^2 \end{cases}$$

where

$$h_1(v_1) = \int_0^{v_1} f(v_1, t) dt = \frac{1}{a^4} \left\{ \pi a^2 - 4av_1^{\frac{1}{2}} + v_1 \right\}, \quad 0 \leq v_1 \leq a^2$$

and

$$b_2(v_1) = \frac{1}{a^4} \left\{ -2a^2 - v_1 + 4a(v_1 - a^2)^{\frac{1}{2}} + 2a^2 \left[\sin^{-1} \frac{a}{\sqrt{v_1}} - \cos^{-1} \frac{a}{\sqrt{v_1}} \right] \right\}, \quad a^2 \leq v_1 \leq 2a^2.$$

Take

$$g_2(v_2) = f_3(v_2) = \frac{1}{a^2} \left\{ av_2^{-\frac{1}{2}} - 1 \right\}, \quad 0 \leq v_2 \leq a^2.$$

Then $u = x^2 = v_1 + v_2$. That part of the joint density of $w_1 = v_1 + v_2$ and $w = v_1$ for $0 \leq v_1 \leq a^2$ and $0 \leq v_2 \leq a^2$ is then given by

$$g_1(w_1, w) = b_1(w)g_2(w_1 - w) = \frac{1}{a^6} \left\{ -\pi a^2 + \pi a^3 (w_1 - w)^{-\frac{1}{2}} + 4aw^{\frac{1}{2}} - 4a^2 w^{\frac{1}{2}} (w_1 - w)^{-\frac{1}{2}} - w + aw(w_1 - w)^{-\frac{1}{2}} \right\}.$$

Note that one has to look into the regions in the (w_1, w) -plane corresponding to the regions $S_1 = \{0 \leq v_1 \leq a^2, 0 \leq v_2 \leq a^2\}$ and $S_2 = \{a^2 \leq v_1 \leq 2a^2, 0 \leq v_2 \leq a^2\}$. For the set S_1 the integral is over $0 \leq w \leq w_1$ when $0 \leq w_1 \leq a^2$ and over $w_1 - a^2 \leq w \leq a^2$ when $a^2 \leq w_1 \leq 2a^2$. Similarly for the set S_2 the integral is when $a^2 \leq w \leq w_1$ when $a^2 \leq w_1 \leq 2a^2$ and over $w_1 - a^2 \leq w \leq 2a^2$ when $2a^2 \leq w_1 \leq 3a^2$. Denoting these four integrals by $f_{1,1}(w_1)$, $f_{1,2}(w_1)$, $f_{2,1}(w_1)$ and $f_{2,2}(w_1)$ respectively one has the following where $f_{1,1}$ and $f_{1,2}$ correspond to S_1 and the remaining to S_2 :

$$\begin{aligned} f_{1,1}(w_1) &= \int_0^{w_1} g_1(w_1, w) dw \\ &= \frac{1}{a^6} \left\{ -3\pi a^2 w_1 + 2\pi a^3 w_1^{\frac{1}{2}} + 4aw_1^{\frac{3}{2}} - \frac{1}{2} w_1^{\frac{3}{2}} \right\}, \quad 0 \leq w_1 \leq a^2 \end{aligned} \quad (2.1)$$

$$\begin{aligned} f_{1,2}(w_1) &= \int_{w_1 - a^2}^{a^2} g_1(w_1, w) dw \\ &= \frac{1}{a^6} \left\{ -2a^2 + (\pi a^2 + a^2)w_1 + \frac{1}{2} w_1^2 - 2\pi a^3 (w_1 - a^2)^{\frac{1}{2}} - 2aw_1 (w_1 - a^2)^{\frac{1}{2}} - 2a(w_1 - a^2)^{\frac{3}{2}} - 4a^2 w_1 \left[\sin^{-1} \frac{a}{\sqrt{w_1}} - \cos^{-1} \frac{a}{\sqrt{w_1}} \right] \right\}, \quad a^2 \leq w_1 \leq 2a^2. \end{aligned} \quad (2.2)$$

The remaining ones $f_{2,1}(w_1)$ and $f_{2,2}(w_1)$ are coming from

$$\begin{aligned}
g_2(w_1, w) &= b_2(w)g_2(w_1 - w), a^2 \leq v_1 \leq 2a^2, 0 \leq v_2 \leq a^2 \\
&= \frac{1}{a^6} \left\{ 2a^2 + w - 2a^3(w_1 - w)^{-\frac{1}{2}} - aw(w_1 - w)^{-\frac{1}{2}} \right. \\
&\quad + 4a^2(w - a^2)^{\frac{1}{2}}(w_1 - w)^{-\frac{1}{2}} - 4a(w - a^2)^{\frac{1}{2}} \\
&\quad + 2a^3(w_1 - w)^{-\frac{1}{2}} \left[\sin^{-1} \frac{a}{\sqrt{w}} - \cos^{-1} \frac{a}{\sqrt{w}} \right] \\
&\quad \left. - 2a^2 \left[\sin^{-1} \frac{a}{\sqrt{w}} - \cos^{-1} \frac{a}{\sqrt{w}} \right] \right\}.
\end{aligned}$$

Then

$$\begin{aligned}
f_{1,2}(w_1) &= \int_{a^2}^{w_1} g_2(w_1, w) dw \\
&= \frac{1}{a^6} \left\{ -\pi a^4 - \frac{5}{2} a^4 + (2\pi a^2 + 2a^2)w_1 + \frac{1}{2} w_1^2 \right. \\
&\quad + (2\pi a^3 - 8a^3)(w_1 - a^2)^{\frac{1}{2}} - 2aw_1(w_1 - a^2)^{\frac{1}{2}} \\
&\quad - 2a(w_1 - a^2)^{-\frac{3}{2}} - 2a^2 w_1 \left[\sin^{-1} \frac{a}{\sqrt{w_1}} - \cos^{-1} \frac{a}{\sqrt{w_1}} \right] \\
&\quad \left. - 8a^3 \int_0^{\cos^{-1} \frac{a}{\sqrt{w_1}}} \left[w_1 - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \right\} \tag{2.3}
\end{aligned}$$

and

$$\begin{aligned}
f_{2,2}(w_1) &= \frac{1}{a^6} \left\{ -\frac{5}{2} a^4 - 3a^2 w_1 - \frac{1}{2} w_1^2 + 8a^3(w_1 - 2a^2)^{\frac{1}{2}} \right. \\
&\quad + 2a(w_1 - 2a^2)^{\frac{3}{2}} + 2aw_1(w_1 - 2a^2)^{\frac{1}{2}} \\
&\quad + 6a^2(w_1 - a^2) \left[\sin^{-1} \frac{a}{\sqrt{w_1 - a^2}} - \cos^{-1} \frac{a}{\sqrt{w_1 - a^2}} \right] \\
&\quad + 4a^2 \left[\sin^{-1} \frac{a}{\sqrt{w_1 - a^2}} - \cos^{-1} \frac{a}{\sqrt{w_1 - a^2}} \right] \\
&\quad \left. - 8a^3 \int_{\cos^{-1} \frac{a}{\sqrt{w_1 - a^2}}}^{\frac{\pi}{4}} \left[w_1 - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \right\}. \tag{2.4}
\end{aligned}$$

Note that $f_{1,2} + f_{2,1}$ is that part of the density for $a^2 \leq w_1 \leq 2a^2$. The density of $u = w_1 = x^2$, denoted by $f(u)$, is the following. This will be stated as a theorem.

Theorem 1.

$$f(u) = \begin{cases} f_{1,1}(u), & 0 \leq u \leq a^2 \\ f_{1,2}(u) + f_{2,1}(u), & a^2 \leq u \leq 2a^2 \\ f_{2,2}(u), & 2a^2 \leq u \leq 3a^2 \end{cases}$$

where $f_{1,1}$ is given in (2.1), $f_{2,1}$ in (2.4) and

$$\begin{aligned} f_{1,2}(u) + f_{2,1}(u) = & \frac{1}{a^6} \left\{ -\pi a^4 - \frac{1}{2} a^4 + (3\pi a^2 + 3a^2)u + u^2 \right. \\ & + 8a^3(u - a^2)^{\frac{3}{2}} - 4au(u - a^2)^{\frac{1}{2}} - 4a(u - a^2)^{\frac{3}{2}} \\ & - 6a^2u \left[\sin^{-1} \frac{a}{\sqrt{u}} - \cos^{-1} \frac{a}{\sqrt{u}} \right] \\ & + 4a^2 \left[\sin^{-1} \frac{a}{\sqrt{w_1 - a^2}} - \cos^{-1} \frac{a}{\sqrt{w_1 - a^2}} \right] \\ & \left. - 8a^3 \int_0^{\cos^{-1} \frac{a}{\sqrt{u}}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \right\}. \end{aligned} \quad (2.5)$$

Then the density of the distance $x = u^{1/2}$, denoted by $g(x)$, is given by

$$g(x) = 2xf(x^2). \quad (2.6)$$

In order to show that $f(u)$ as given above is in fact a density one has to evaluate the two integrals appearing in (2.4) and (2.5). Making the transformation $z = a^2/\cos^2 \theta$ one has

$$I_1 = -8a^3 \int_0^{\cos^{-1} \frac{a}{\sqrt{u}}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta = -4a^4 \int_{a^2}^u \frac{(u-z)^{\frac{1}{2}}}{z\sqrt{z-a^2}} dz, \quad a^2 \leq u \leq 2a^2 \quad (2.7)$$

and

$$\begin{aligned} I_2 = & -8a^3 \int_{\cos^{-1} \frac{a}{\sqrt{u-a^2}}}^{\frac{\pi}{4}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \\ = & -4a^4 \int_{u-a^2}^{2a^2} \frac{(u-z)^{\frac{1}{2}}}{z\sqrt{z-a^2}} dz, \quad 2a^2 \leq u \leq 3a^2. \end{aligned} \quad (2.8)$$

Combining the regions $\{a^2 \leq u \leq 2a^2, a^2 \leq z \leq u\}$ and $\{2a^2 \leq u \leq 3a^2, u - a^2 \leq z \leq 2a^2\}$ and then changing the order of integration the sum of the two integrals can be evaluated. Since the steps in this derivation are going to be used later on, this result will be stated as a lemma.

Lemma 1.

$$\begin{aligned} I &= \int_{a^2}^{2a^2} I_1 du + \int_{2a^2}^{3a^2} I_2 du = -4a^4 \left\{ \int_{z=a^2}^{2a^2} \frac{1}{z\sqrt{z-a^2}} \left[\int_{u=z}^{z+a^2} (u-z)^{\frac{1}{2}} du \right] dz \right\} \\ &= -\frac{8}{3} a^7 \int_{z=a^2}^{2a^2} \frac{1}{z\sqrt{z-a^2}} dz = -\frac{8}{3} a^7 \int_{z=0}^{a^2} \frac{1}{(z-a^2)\sqrt{z}} dz = -\frac{1}{3} \pi a^6 \end{aligned} \quad (2.9)$$

by making the substitution $z = a^2 \tan^2 \phi$.

The general b -th moment of u_1 that is $E(u^b) = E(x^{2b})$, for an arbitrary b , can be evaluated by computing the integrals

$$\int_0^{a^2} u^b f_{1.1}(u) du, \int_{a^2}^{2a^2} u^b (f_{1.2} + f_{2.1}) du, \int_{2a^2}^{3a^2} u^b f_{2.2}(u) du.$$

In order to achieve this we need some integrals which will be stated as lemmas here. Special cases for $b = 1/2, 1$ are needed if we want to obtain the mean value and variance of x and hence these special cases are also listed with some of the lemmas.

Lemma 2. For $0 < \gamma \leq 1$, $\Re(b) > -1$, $\Re(\delta) > -1$, where $\Re(\cdot)$ denotes the real part of (\cdot)

$$\begin{aligned} \int_0^\gamma v^\delta (1+v)^b dv &= \frac{1}{(\delta+1)^2} F_1(-b, \delta+1; \delta+2; -\gamma) \\ &= \frac{\sqrt{3}}{4} - \frac{1}{4} \ln \sqrt{2} + \frac{1}{4} \ln(\sqrt{3}-1) \text{ for } \delta = \frac{1}{2}, b = \frac{1}{2}, \gamma = \frac{1}{2} \\ &= \frac{1}{8} \ln \sqrt{2} - \frac{1}{8} \ln(\sqrt{3}-1) \text{ for } \delta = \frac{3}{2}, b = \frac{1}{2}, \gamma = \frac{1}{2} \\ &= \frac{3}{4} \sqrt{2} - \frac{1}{4} \ln(\sqrt{2}-1) \text{ for } \delta = \frac{1}{2}, b = \frac{1}{2}, \gamma = 1 \\ &= \frac{7}{24} \sqrt{2} - \frac{1}{8} \ln(\sqrt{2}-1) \text{ for } \delta = \frac{3}{2}, b = \frac{1}{2}, \gamma = 1 \\ &= \frac{19\sqrt{2}}{280} \text{ for } \delta = \frac{3}{2}, b = 1, \gamma = 1 \\ &= \frac{13\sqrt{2}}{60} \text{ for } \delta = \frac{1}{2}, b = 1, \gamma = \frac{1}{2} \\ &= \frac{24}{35} \text{ for } \delta = \frac{3}{2}, b = 1, \gamma = 1 \\ &= \frac{16}{15} \text{ for } \delta = \frac{1}{2}, b = 1, \gamma = 1, \end{aligned}$$

where ${}_2F_1(\cdot)$ is a Gauss' hypergeometric function.

The integral is evaluated by expanding $(1+v)^b$ and integrating term by term.

Lemma 3. For $\gamma > 1$, $\Re(\alpha) > -1$

$$\int_0^{\frac{1}{\gamma}} t^\alpha (1-t)^{-\beta} (\gamma-t)^{-\delta} dt = \frac{\gamma^{-\delta-1}}{(\alpha+1)} F_1\left(\alpha+1, \beta, \delta; \alpha+2; \frac{1}{\gamma}, \frac{1}{\gamma^2}\right)$$

where F_1 is an Appell's double hypergeometric function which is also the Lauricella function F_D for two variables, see for example Mathai (1993).

The integral is evaluated by expanding $(1-t)^{-\beta}$ and $\left(1-\frac{t}{\gamma}\right)^{-\delta}$ by using binomial expansions, integrating out and then interpreting the double series.

Lemma 4. For $m = 1, 2, \dots$

$$\begin{aligned} \int \frac{1}{(\cos \theta)^{2m}} d\theta &= \frac{\sin \theta}{(2m-1)} \left\{ \frac{1}{(\cos \theta)^{2m-1}} + \frac{2(m-1)}{(2m-3)} \frac{1}{(\cos \theta)^{2m-3}} \right. \\ &+ \frac{2^2(m-1)(m-2)}{(2m-3)(2m-5)} \frac{1}{(\cos \theta)^{2m-5}} + \dots \\ &\left. + 2^{m-1} \frac{(m-1)(m-2)\dots 1}{(2m-3)(2m-5)\dots 1} \frac{1}{\cos \theta} \right\} \end{aligned}$$

and

$$\begin{aligned} \int \frac{1}{(\cos \theta)^{2m+1}} d\theta &= \frac{\sin \theta}{2m} \left\{ \frac{1}{(\cos \theta)^{2m}} \right. \\ &+ \sum_{k=1}^{m-1} \frac{(2m-1)(2m-3)\dots(2m-2k+1)}{2^k(m-1)(m-2)\dots(m-k)} \frac{1}{(\cos \theta)^{2m-2k}} \\ &\left. + \frac{(2m-1)!}{2^m m!} \ln\left(\frac{\cos \theta}{1-\sin \theta}\right) \right\}. \end{aligned}$$

The results are available by successive integration, see also Gradshteyn and Ryzhik (1980) formulae 2.519.1 and 2.519.2. But for computing arbitrary moments of x or of $u = x^2$ we need $E(u^b)$ for an arbitrary b which in turn will require $\int (\cos^2 \theta)^{-\alpha} d\theta$ for an arbitrary α . This can be by substituting $t = \sin \theta$, expanding $(1-t^2)^{-(\alpha+1/2)}$ with the help of a binomial expansion and then integrating. The final result is stated next.

Lemma 5.

$$\int \frac{1}{(\cos^2 \theta)^\alpha} d\theta = \sin \theta {}_2F_1\left(\alpha + \frac{1}{2}, \frac{1}{2}; \frac{3}{2}; \sin^2 \theta\right).$$

It can be seen that for $\alpha = 1, 2, \dots$ this result agrees with those in Lemma 4. In order to see this, use the result

$${}_2F_1(a, b; c; x) = (1-x)^{-b} {}_2F_1(c-a, b; c; x/(x-1)).$$

With the help of the above lemmas the b -th moment of u , for an arbitrary b , is the following:

Theorem 2. For $\Re(b) > -1$,

$$\begin{aligned} E(u^b) = & a^{2b+6} \left\{ \frac{b_1}{(b+1)} + \frac{b_2}{(b+2)} + \frac{b_3}{(b+3)} \right. \\ & + \frac{b_4}{(2b+3)} + \frac{b_5}{(2b+4)} + \frac{b_6}{(2b+5)} + \phi_1 + \phi_2 \\ & - \frac{16}{3} {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -1\right) - \frac{8}{3} {}_2F_1\left(-b-1, \frac{3}{2}; \frac{5}{2}; -1\right) \\ & - \frac{8}{5} {}_2F_1\left(-b, \frac{5}{2}; \frac{7}{2}; -1\right) + \frac{16}{3} 2^b {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -\frac{1}{2}\right) \\ & + \frac{4}{5} 2^b {}_2F_1\left(-b, \frac{5}{2}; \frac{7}{2}; -\frac{1}{2}\right) \\ & + \frac{2^{b+4}\sqrt{2}}{(b+1)} F_1\left(\frac{1}{2}, b+\frac{3}{2}, -b-1; \frac{3}{2}; \frac{1}{2}, \frac{1}{4}\right) \\ & + 3\pi 2^{b-1} F_1\left(1, -b, b+3; 2; \frac{1}{4}, \frac{1}{2}\right) \\ & + 12 \frac{2^{b+1}\sqrt{2}}{(b+2)} F_1\left(\frac{1}{2}, b+\frac{5}{2}, -b-2; \frac{3}{2}; \frac{1}{2}, \frac{1}{4}\right) \\ & \left. - 12 \frac{2^b\sqrt{2}}{(b+1)} F_1\left(\frac{1}{2}, b+\frac{3}{2}, -b-1; \frac{3}{2}; \frac{1}{2}, \frac{1}{4}\right)\right\} \end{aligned}$$

where

$$b_1 = \pi + \frac{1}{2} + (2-3\pi)2^{b+1} + \left(3\pi - \frac{5}{2}\right)3^{b+1}$$

$$b_2 = -3(1+2\pi) + 3(2+\pi)2^{b+2} - 3(1+\pi)3^{b+2}$$

$$b_3 = -\frac{3}{2} + \frac{3}{2}2^{b+3} - \frac{1}{2}3^{b+3}$$

$$b_4 = 4\pi, \quad b_5 = 6\pi, \quad b_6 = 8$$

$$\phi_1 = -\frac{24}{(2b+4)} \int_0^{\frac{\pi}{4}} \frac{1}{(\cos \theta)^{2b+4}} d\theta$$

and

$$\begin{aligned}
\phi_2 &= -8a^3 \left\{ \int_{u=a^2}^{2a^2} u^b \int_0^{\cos^{-1} \frac{a}{\sqrt{u}}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \right. \\
&\quad \left. + \int_{u=2a^2}^{3a^2} u^b \int_{\cos^{-1} \frac{a}{\sqrt{u-a^2}}}^{\frac{\pi}{4}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \right\} \\
&= -4a^2 \left\{ \int_{z=a^2}^{2a^2} \frac{1}{z\sqrt{z-a^2}} \left[\int_{u=z}^{z+a^2} u^b (u-z)^{\frac{1}{2}} du \right] dz \right\} \tag{2.10}
\end{aligned}$$

by using Lemma 1.

The last integral above can be written as follows:

$$\int_{u=z}^{z+a^2} u^b (u-z)^{\frac{1}{2}} du = 2a^{b+\frac{3}{2}} \int_0^{\tan^{-1} \frac{a}{\sqrt{z}}} \left[\frac{1}{(\cos^2 \theta)^{b+z}} - \frac{1}{(\cos^2 \theta)^{b+1}} \right] d\theta.$$

Now, with the help of Lemmas 4 and 5, ϕ_1 and ϕ_2 can be written explicitly.

3. ONE POINT ON A FACE AND THE OTHER POINT INSIDE

Let $z_2 = 0$. In this case the point P is inside the cube and the point Q is on a face. Taking this face as the (x, y) -plane and referring back to the general procedure we have the following: $g_1(v_1)$ remains the same and

$$g_2(v_2) = \frac{v_2^{-\frac{1}{2}}}{2a}, \quad 0 \leq v_2 \leq a^2.$$

Then proceeding as in the general case the density of $u = x^2$, denoted by $f^{(1)}(u)$, is the following:

Theorem 3.

$$f^{(1)}(u) = \frac{1}{a^5} \begin{cases} \rho_1(u), & 0 \leq u \leq a^2 \\ \rho_2(u), & a^2 \leq u \leq 2a^2 \\ \rho_3(u), & 2a^2 \leq u \leq 3a^2 \end{cases}$$

where

$$\rho_1(u) = \pi a^2 u^{\frac{1}{2}} - \pi a u + \frac{2}{3} u^{\frac{3}{2}},$$

$$\rho_1(u) = -\frac{1}{3} a^3 + (a + \pi a)u - 2a^2(u - a^2)^{\frac{1}{2}} - 2u(u - a^2)^{\frac{1}{2}} + \frac{2}{3}(u - a^2)^{\frac{3}{2}} \\ - 2au \left[\sin^{-1} \frac{a}{\sqrt{u}} - \cos^{-1} \frac{a}{\sqrt{u}} \right] - 4a^2 \int_0^{\cos^{-1} \frac{a}{\sqrt{u}}} \left[u - \frac{a^2}{\cos^2 \theta} \right] d\theta$$

and

$$\rho_3(u) = -\frac{5}{3} a^3 - au + 2a^2(u - 2a^2)^{\frac{1}{2}} + u(u - 2a^2)^{\frac{1}{2}} \\ = -\frac{1}{3}(u - 2a^2)^{\frac{3}{2}} + 2a^3 \left[\sin^{-1} \frac{a}{\sqrt{u - a^2}} - \cos^{-1} \frac{a}{\sqrt{u - a^2}} \right] \\ - 2a(u - a^2) \left[\sin^{-1} \frac{a}{\sqrt{u - a^2}} - \cos^{-1} \frac{a}{\sqrt{u - a^2}} \right] \\ - 4a^2 \int_{\cos^{-1} \frac{a}{\sqrt{u - a^2}}}^{\frac{\pi}{4}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta.$$

The general b -th moment, for an arbitrary b , can be computed by integrating with the help of Lemmas 1 to 5. The final result is the following, the details of the steps are omitted for saving space:

Theorem 4. For $\Re(b) > -1$,

$$\mathbb{E}(u^b) = a^{2b} \left\{ \frac{2\pi}{(2b+3)} + \frac{4}{3(2b+5)} + \frac{1}{(b+1)} \left[\frac{1}{3} + \frac{2^{b+3}}{3} - 5(3^b) \right] \right. \\ + \frac{1}{(b+2)} [-1 - \pi + 2^{b+3} - 3^{b+2}] \\ - \frac{4}{3} {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -1\right) - \frac{4}{3} {}_2F_1\left(-b, -1, \frac{3}{2}; \frac{5}{2}; -1\right) \\ + \frac{4}{15} {}_2F_1\left(-b, \frac{5}{2}; \frac{7}{2}; -1\right) - \frac{2\sqrt{2}}{(b+2)} {}_2F_1\left(b + \frac{5}{2}, \frac{1}{2}; \frac{3}{2}; \frac{1}{2}\right) \\ + \frac{2^{b+2}}{3} {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -\frac{1}{2}\right) + \frac{2^{b+2}}{3} {}_2F_1\left(-b, -1, \frac{3}{2}; \frac{5}{2}; -\frac{1}{2}\right) \\ - \frac{2^{b+1}}{15} {}_2F_1\left(-b, \frac{5}{2}; \frac{7}{2}; -\frac{1}{2}\right) \\ \left. + \frac{2^{b+3}\sqrt{2}}{(b+2)} F_1\left(\frac{1}{2}, b + \frac{5}{2}, -b - 2; \frac{3}{2}; \frac{1}{2}, \frac{1}{4}\right) + \psi \right\}$$

where

$$\begin{aligned} \psi &= -4a^2 \int_{a^2}^{2a^2} u^b \left\{ \int_0^{\cos^{-1} \frac{a}{\sqrt{u}}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \right. \\ &\quad \left. - 4a^2 \int_{2a^2}^{3a^2} u^b \left\{ \int_{\frac{\pi}{4}}^{\cos^{-1} \frac{a}{\sqrt{u-a^2}}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \right\} d\theta \right. \\ &= -2a^3 \int_{z=a^2}^{2a^2} \frac{1}{z\sqrt{z-a^2}} \left[\int_{u=z}^{z+a^2} u^b (u-z)^{\frac{1}{2}} du \right] dz \end{aligned}$$

by using Lemma 1,

$$\begin{aligned} &= -2a^3 \int_{z=a^2}^{2a^2} \frac{1}{z\sqrt{z-a^2}} \left[\int_{v=0}^{a^2} v^{\frac{1}{2}} (v-z)^b du \right] dz \\ &\quad - 4a^3 \int_{z=a^2}^{2a^2} u^b \left[\int_0^{\tan^{-1} \frac{a}{\sqrt{z}}} \left\{ \frac{1}{(\cos^2 \phi)^{b+2}} - \frac{1}{(\cos^2 \phi)^{b+1}} \right\} d\phi \right] dz. \end{aligned}$$

Since this integral goes into a multiple series for general values of b no further rewriting will be attempted here.

4. THE DISTANCE BETWEEN TWO RANDOM POINTS ON OPPOSITE SIDES OF A CUBE

Let the point P be on a given face. Take one corner of this face as the origin of a rectangular coordinate system and this face on the (x, y) - plane. Let the point Q be on the opposite face. Then the coordinates of P and Q can be written as $P = P(x_1, y_1, 0)$, $Q = Q(x_2, y_2, a)$. The squared distance between P and Q is the given by

$$u = (x_2 - x_1)^2 + (y_2 - y_1)^2 + a^2.$$

When a point is uniformly distributed over the surface of a cube of side a the probability that the point is on a given side is $\frac{a^2}{6a^2} = \frac{1}{6}$ and this probability remains the same for all the six faces. Hence, due to symmetry we need to consider only the situation where P is uniformly distributed over the (x, y) - face and Q is independently and uniformly distributed over the opposite face. In other words, we need to assume only that $x_j, y_j, j = 1, 2$ are mutually independently and uniformly distributed over $[0, a]$. Note that $u = v_1 + a^2$ where the density of v_1 is $b_2(v_1)$, and from there we have the density of u which is denoted by $f^{(2)}(u)$ in the following theorem.

Theorem 5.

$$f^{(2)}(u) = \frac{1}{a^4} \begin{cases} \psi_1(u), & a^2 \leq u \leq 2a^2 \\ \psi_2(u), & 2a^2 \leq u \leq 3a^2 \end{cases} \quad (4.1)$$

where

$$\psi_1(u) = \pi a^2 - 4a(u - a^2)^{\frac{1}{2}} + (u - a^2)$$

and

$$\psi_2(u) = -2a^2 - (u - a^2) + 4a(u - 2a^2)^{\frac{1}{2}} + 2a^2 \left[\sin^{-1} \frac{a}{\sqrt{u - a^2}} - \cos^{-1} \frac{a}{\sqrt{u - a^2}} \right].$$

4.1. The general b -th moment of u

Evaluation of the positive integer moments of u is not difficult. But if positive integer moments of $x = u^{1/2}$ are required then one has to consider positive integer and half-integer values of b . For computing the b -th moment of u , for an arbitrary b , one needs the help of the lemmas given earlier. The following is the final result.

Theorem 6. For $\Re(b) > -1$,

$$\begin{aligned} \mathbb{E}(u^b) &= a^{2b} \left\{ \frac{1}{(b+1)} [-\pi + 1 - 3^{b+1}] + \frac{1}{(b+1)} [2^{b+3} - 3^{b+2} - 1] \right. \\ &\quad - \frac{8}{3} {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -1\right) + \frac{2^{b+3}}{3} {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -\frac{1}{2}\right) \\ &\quad \left. + \sqrt{2} \frac{2^{\frac{b+5}{2}}}{(b+1)} F_1\left(\frac{1}{2}, -b-1, b+\frac{3}{2}; \frac{3}{2}; \frac{1}{4}, \frac{1}{2}\right) \right\}. \end{aligned} \quad (4.2)$$

Substituting $b = 1/2$, $b = 1$ one has the following expressions for the expected values of x and x^2 respectively: From there, the variance of x is readily available

$$\begin{aligned} \mathbb{E}(x) &= \mathbb{E}(u^{\frac{1}{2}}) = a \left\{ -\frac{2}{3} \pi + \frac{\sqrt{2}}{5} - \frac{8}{5} \sqrt{3} + \frac{4}{15} - 4 \ln \sqrt{2} - \ln(\sqrt{2} - 1) \right. \\ &\quad \left. + 4 \ln(\sqrt{3} - 1) + \frac{16}{3} F_1\left(\frac{1}{2}, -\frac{3}{2}, 2; \frac{3}{2}; \frac{1}{4}, \frac{1}{2}\right) \right\} \end{aligned} \quad (4.3)$$

and

$$\mathbb{E}(x^2) = \mathbb{E}(u) = \frac{4}{3} a^2. \quad (4.4)$$

4.2. Distance between two random points in a square

Observe that v_1 is the squared distance between two independently and uniformly distributed random points in a square. Arbitrary moments of this squared distance in a square is then $E(v_1^b)$ for an arbitrary b , which is available by direct integration. The result is the following:

Corollary 4.1. For $\Re(b) > -1$

$$E(v_1^b) = a^{2b} \left\{ 2(1 - 2^{b+1}) \left[\frac{1}{(b+1)} + \frac{1}{(b+2)} \right] - \frac{8}{(2b+3)} \right. \\ \left. + \frac{8}{3} {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -1\right) + \frac{2\sqrt{2}}{(b+1)} {}_2F_1\left(b + \frac{3}{2}, \frac{1}{2}; \frac{3}{2}; \frac{1}{2}\right) \right\}. \quad (4.5)$$

5. RANDOM POINTS ON ADJACENT FACES

Put $y_2 = a$, $z_1 = 0$. Then the point P is on the (x, y) - plane and the other point Q is on the face $y_2 = a$. Since y_i and $a - y_i$ are identically distributed, and the same goes for x_i as well as z_i , this case is the same as saying that the two points are on adjacent faces. The distance between P and Q in this case is also the length of the secant across two adjacent faces or random path across two adjacent faces when the points are independently and uniformly distributed over the respective faces.

Referring back to our notations we have

$$f_1(u_1) = \frac{1}{a^2} \left\{ au_1^{-\frac{1}{2}} - 1 \right\}, \quad 0 \leq u_1 \leq a^2$$

$$f_2(u_2) = \frac{u_2^{-\frac{1}{2}}}{2a}, \quad 0 \leq u_2 \leq a^2$$

and let

$$f_3(u_3) = \frac{u_3^{-\frac{1}{2}}}{2a}, \quad 0 \leq u_3 \leq a^2.$$

Then proceeding as before the various expressions are the following:

$$g_1(v_1) = \frac{1}{a^3} \begin{cases} \frac{\pi a}{2} - v_1^{\frac{1}{2}}, & 0 \leq v_1 \leq a^2 \\ -a + (v_1 - a^2)^{\frac{1}{2}} + a \left[\sin^{-1} \frac{a}{\sqrt{v_1}} - \cos^{-1} \frac{a}{\sqrt{v_1}} \right], & a^2 \leq v_1 \leq 2a^2 \end{cases}$$

and let

$$g_2(v_2) = f_3(v_2).$$

Consequently, the density of $u = \kappa^2$, where κ is the distance between P and Q , denoted by $f^{(3)}(u)$, is the following:

Theorem 7.

$$f^{(3)}(u) = \frac{1}{a^4} \begin{cases} \eta_1(u), & 0 \leq u \leq a^2 \\ \eta_2(u), & a^2 \leq u \leq 2a^2 \\ \eta_3(u), & 2a^2 \leq u \leq 3a^2 \end{cases} \quad (5.1)$$

where

$$\eta_1(u) = \frac{\pi a}{2} u^{\frac{1}{2}} - \frac{\pi}{4} u,$$

$$\begin{aligned} \eta_2(u) = & \frac{\pi a^2}{4} + \frac{\pi}{4} u - a(u - a^2)^{\frac{1}{2}} - \frac{u}{2} \left[\sin^{-1} \frac{a}{\sqrt{u}} - \cos^{-1} \frac{a}{\sqrt{u}} \right] \\ & - 2a \int_0^{\cos^{-1} \frac{a}{\sqrt{u}}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta \end{aligned}$$

and

$$\begin{aligned} \eta_3(u) = & -a^2 + a(u - 2a^2)^{\frac{1}{2}} \\ & + \frac{1}{2}(u - a^2) \left[\sin^{-1} \frac{a}{\sqrt{u - a^2}} - \cos^{-1} \frac{a}{\sqrt{u - a^2}} \right] \\ & + a^2 \left[\sin^{-1} \frac{a}{\sqrt{u - a^2}} - \cos^{-1} \frac{a}{\sqrt{u - a^2}} \right] \\ & - 2a \int_{\cos^{-1} \frac{a}{\sqrt{u - a^2}}}^{\frac{\pi}{4}} \left[u - \frac{a^2}{\cos^2 \theta} \right]^{\frac{1}{2}} d\theta. \end{aligned}$$

In order to see that $f^{(3)}(u)$ is in fact a density we need to evaluate the integrals in $\eta_2(u)$ and $\eta_3(u)$. The steps in this procedure are also needed when computing the arbitrary moments u in this case. This is done with the help of Lemma 1 and the final result is the following:

Theorem 8. For $\Re(b) > -1$

$$\begin{aligned}
 E(u^b) = a^{2b} & \left\{ \frac{1}{(b+1)} \left[2^{b+1} - 3^{b+1} - \frac{\pi}{4} \right] - \frac{\pi}{4(b+2)} + \frac{\pi}{(2b+3)} \right. \\
 & + \frac{2^{b+1}}{3} {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -\frac{1}{2}\right) - \frac{2}{3} {}_2F_1\left(-b, \frac{3}{2}; \frac{5}{2}; -1\right) \\
 & - \frac{1}{(b+2)} \frac{1}{\sqrt{2}} {}_2F_1\left(-b + \frac{5}{2}, \frac{1}{2}; \frac{3}{2}; \frac{1}{2}\right) \\
 & + \frac{2^{\frac{b+3}{2}}}{(b+1)} F_1\left(\frac{1}{2}, -b-1, b + \frac{3}{2}; \frac{3}{2}; \frac{1}{4}, \frac{1}{2}\right) \\
 & + \frac{2^{\frac{b+3}{2}}}{(b+2)} F_1\left(\frac{1}{2}, -b-2, b + \frac{5}{2}; \frac{3}{2}; \frac{1}{4}, \frac{1}{2}\right) \\
 & \left. - \frac{2^{\frac{b+1}{2}}}{(b+1)} F_1\left(\frac{1}{2}, -b-1, b + \frac{3}{2}; \frac{3}{2}; \frac{3}{2}; \frac{1}{4}, \frac{1}{2}\right) + \phi \right\} \quad (5.2)
 \end{aligned}$$

where ϕ is coming from the last two integrals in $f^{(3)}(u)$, which is given below.

$$\phi = -a^2 \left\{ \int_{z=a^2}^{2a^2} \frac{1}{z\sqrt{z-a}} \left[\int_{u=z}^{z+a^2} u^b (u-z)^{\frac{1}{2}} du \right] dz \right\}. \quad (5.3)$$

Put $v = u - z$, $z = a^2 t$, $v = a^2 w$ to rewrite




$$\phi = -a^{2b+4} \left\{ \int_0^1 w^{\frac{1}{2}} \left[\int_0^1 \frac{(t+(1+w))^b}{\sqrt{t}(1+t)} dt \right] dw \right\}. \quad (5.4)$$





For positive integer values of b this double integral is easily evaluated either by expanding $(t+(1+w))^b$ with the help of a binomial expansion and then integrating out or by starting with (5.3), writing the inner integral as $\int_0^{a^2} v^{1/2} (v+z)^b dv$, expanding $(v+z)^b$ and then integrating out. But for arbitrary b it does not seem that ϕ can be put into any simple form. One can write ϕ in terms of a general triple hypergeometric series. This will be given here in order to save space.




6. GRAPHS, SIMULATION RESULTS AND REMARKS


The following are the graphs of the densities, drawn by using a mathematica program, showing the behaviour of the tail and mid-regions along with the combined graphs for a cube of side 1, that is $a = 1$.

The simplest case is the density in Theorem 5 which behaves like an exponential curve. The next ones correspond to the densities in Theorem 7, Theorem 3 and Theorem 1 respectively. The curves seem to approximate progres-












The density in Theorem 7 for $\alpha=1$, left tail



The density in Theorem 7 for $\alpha=1$, middle portion



The density in Theorem 7 for $\alpha=1$, right tail



The density in Theorem 7 for $\alpha=1$

sively to that of a gamma type density. The right tail contributes less and less as one goes from Theorem 7 to Theorem 3 to Theorem 1. Fairly good approximations to the densities are available by deleting fully the right tail terms in Theorem 3 and 1 and the terms containing the integrals in Theorem 7.

Department of Mathematics and Statistics
McGill University, Canada

A.M. MATHAI

Department of Mathematical Sciences
The University of Texas at El Paso, U.S.A

PETER MOSCHOPOULOS

Department of Quantitative Methods
Libero Istituto Universitario Carlo Cattaneo, Italy

GIORGIO PEDERZOLI

REFERENCES

- R. COLEMAN (1969), *Random path through convex bodies*, "Journal of Applied Probability", 6, pp. 430-441.
- S. EILON, C.D.T. WATSON-GANDY, N. CHRISTOFIDES (1971), *Distribution management: mathematical modelling and practical analysis*, Griffin, London.
- B. GHOSH (1943a), *On the distribution of random distances in a rectangle*, "Science and Culture", 8(9), pp. 388.
- B. GHOSH (1943b), *On random distances between two rectangles*, "Science and Culture", 8(11), pp. 464.
- B. GHOSH (1949), *Topografic variation in statistical fields*, "Bulletin of Calcutta Statistical Association", 2(5), pp. 11-28.
- B. GHOSH (1951), *Random distances within a rectangle and between two rectangles*, "Bulletin of Calcutta Statistical Association", 43, pp. 17-24.
- I.S. GRADSHTEYN, I.M. RYZHIK (1980), *Tables of integrals, series, and products*, Academic Press, New York.
- M. HOROWITZ (1965), *Probability of random paths across elementary geometrical shapes*, "Journal of Applied Probability", 2, pp. 169-177.
- G. JÄGER (1911), *Zur theorie des nachbells*, "Sitzungsberichte, Wiener Akademie, Mathematische-Naturwissenschaften Klasse", 120, II, pp. 613-634.
- M.G. KENDALL, P.A.P. MORAN (1963), *Geometrical probability*, Griffin, London.
- J.F.C. KINGMAN (1965), *Mean free paths in a convex reflecting region*, "Journal of Applied Probability", 2, pp. 162-168.
- J.F.C. KINGMAN (1969), *Random secants of a convex body*, "Journal of Applied Probability", 6, pp. 660-672.
- A.M. MATHAI (1993), *A handbook of generalized special functions for statistical and physical sciences*, Oxford University Press, Oxford.
- P.A.P. MORAN (1966), *A note on recent research in geometrical probability*, "Journal of Applied Probability", 3, pp. 453-463.
- P.A.P. MORAN (1969), *A second note on recent research in geometrical probability*, "Advances in Applied Probability", 1, pp. 73-89.
- O. PERSSON (1964), *Distance methods*, "Studia Forestalia Suecica", 15.
- O. PERSSON (1965), *Distance methods*, "Research Note No. 6", Department of Forest Biometry, Royal College of Forestry, Stockholm.

R.J. SMEED, G.O. JEFFCOATE (1963), *Traffic flow during the journey to work in the central area of a town which has a rectangular grid for its road system*, "Proceedings of the Second International Symposium on the Theory of Road Traffic Flow", J. Almond ed., OECD, London, pp. 369-390.

RIASSUNTO

Distanze tra punti aleatori di un cubo

Vengono fornite le forme esplicite delle densità e dei momenti relativi alla distanza tra due punti aleatori distribuiti in modo uniforme ed indipendente all'interno di un cubo. Casi particolari riguardano situazioni dove i punti si trovano su facce opposte oppure adiacenti. Espressioni esatte per momenti arbitrari sono ottenute in termini di funzioni di Gauss, Appell e Lauricella.

SUMMARY

Distance between random points in a cube

The explicit forms of the exact density and the exact arbitrary moments of the distance between two random points when the points are independently and uniformly distributed inside a cube are given. Special cases considered include the density and arbitrary moments of the distance between two independent random points where one is inside the cube and the other is on a face, points on adjacent faces, points on opposite faces and random points inside a square. Exact arbitrary moments are obtained in terms of Gauss' hypergeometric functions and Appell's functions or Lauricella functions.

BIOMETRICS

JOURNAL OF THE BIOMETRIC SOCIETY

Vol. 54, No. 4

December 1998

CONTENTS

- D.L. Borchers, W. Zucchini, R.M. Fewster: Mark-Recapture Models for Line Transect Surveys.
D.L. Borchers, S.T. Buckland, P.W. Goedhart, E.D. Clarke, S.L. Hedley: Horvitz-Thompson Estimators for Double-Platform Line Transect Surveys.
E.R. Hauser, M. Boehnke: Genetic Linkage Analysis of Complex Genetic Traits by Using Affected Sibling Pairs.
J. Teng, D. Siegmund: Multipoint Linkage Analysis Using Affected Relative Pairs and Partially Informative Markers.
J.M. Olson: Discussion of Teng and Siegmund Article.
D.L. Nicolae, M.L. Frigge, N.J. Cox, A. Kong: Discussion.
J. Teng, D. Siegmund: Rejoinder.
L. Hiby, P. Lovell: Using Aircraft in Tandem Formation to Estimate Abundance of Harbour Porpoise.
K.B. Newman: State-Space Modeling of Animal Movement and Mortality with Application to Salmon.
F.A. van Eeuwijk, P.M. Kroonenberg: Multiplicative Models for Interaction in Three-Way ANOVA, with Applications to Plant Breeding.
D.A. Stephens, R.D. Fisch: Bayesian Analysis of Quantitative Trait Locus Data Using Reversible Jump Markov Chain Monte Carlo.
M.-H. Chen, A.K. Manatunga, C.J. Williams: Heritability Estimates from Human Twin Data by Incorporating Historical Prior Information.
D.J. Young, M. Bbandary: Test for Equality of Intraclass Correlation Coefficients Under Unequal Family Sizes.
J.-M. Azaïs, H. Monod, R.A. Bailey: The Influence of Design on Validity and Efficiency of Neighbour Methods.
Y.-C. Huang, W.-K. Wong: Sequential Construction of Multiple-Objective Optimal Designs.
Y. Mehrabi, J.N.S. Matthews: Implementable Bayesian Designs for Limiting Dilution Assays.
P. Hu, A.A. Tsiatis, M. Davidian: Estimating the Parameters in the Cox Model When Covariate Variables Are Measured with Error.
X. Huang, S. Chen, S.-J. Soong: Piecewise Exponential Survival Trees with Time-Dependent Covariates.
H.T. Kim, Y.K. Truong: Nonparametric Regression Estimates with Censored Data: Local Linear Smoothers and Their Applications.
U.G. Dafni, A.A. Tsiatis: Evaluating Surrogate Markers of Clinical Outcome When Measured with Error.
D. Sinha: Posterior Likelihood Methods for Multivariate Survival Data.
D. Faraggi, R. Simon: Bayesian Variable Selection Method for Censored Survival Data.
D.J. Sargent: A General Framework for Random Effects Survival Analysis in the Cox Proportional Hazards Setting.
W.B. Goggins, D.M. Finkelstein, D.A. Schoenfeld, A.M. Zaslavsky: A Markov Chain Monte Carlo EM Algorithm for Analyzing Interval-Censored Data Under the Cox Proportional Hazards Model.
A. Trodikov: A Proportional Hazards Model Taking Account of Long-Term Survivors.
P.J.M. Verweij, H.C. van Houwelingen, T. Stijnen: A Goodness-of-Fit Test for Cox's Proportional Hazards Model Based on Martingale Residuals.
A.B. Anderson, R.L. Prentice: On the Accommodation of Disease Rate Correlations in Aggregate Data Studies of Disease Risk Factors.
V.W. Berger, T. Permutt, A. Ivanova: Convex Hull Test for Ordered Categorical Data.
P.A. Flyer: A Comparison of Conditional and Unconditional Randomization Tests for Highly Stratified Designs.
C.H. Morrel: Likelihood Ratio Testing of Variance Components in the Linear Mixed-Effects Model Using Restricted Maximum Likelihood.
S.G. Baker: Evaluating the Age to Begin Periodic Breast Cancer Screening Using Data from a Few Regularly Scheduled Screenings.
T. Park: An Approach to Categorical Data with Nonignorable Nonresponse.
D.J. Causeur, T.J. Dornbe: Finite Sample Properties of a Multivariate Extension of Double Regression.
R.W. Falk, G.G. Koch: Testing a Specified Difference Between Proportions.

SHORTER COMMUNICATIONS

- C.R. Mehta, N.R. Patel, P. Senchaudhuri: Exact Power and Sample-Size Computations for the Cochran-Armitage Trend Test.
J. Shults, N. Rao Chaganty: Analysis of Serially Correlated Data Using Quasi-Least Squares.
X. Xue: Multivariate Survival Data Under Bivariate Frailty: An Estimating Equation Approach.
E.W. Lee, M.Y. Kim: The Analysis of Correlated Panel Data Using a Continuous-Time Markov Model.
T. Park, D.W. Shin, C.G. Park: A Generalized Estimating Equations Approach for Testing Ordered Group Effects with Repeated Measurements.

CONSULTANT'S FORUM

- J.N. Variyam, J. Blaylock, D. Smallwood: Reduced-Rank Models for Nutrition Knowledge Assessment.
M. Lunn: Applying k -Sample Tests to Conditional Probabilities for Competing Risks in a Clinical Trial.

Correspondence - Book Reviews.
