

Last class:

$$y' = f(t, y)$$

$$y(0) = y_0$$

formula $y(t)$

2 methods to analytically solve the ODE.

if $f(t, y) = g(t) h(y)$


then, $\int \frac{dy}{h(y)} = \int g(t) dt$

Tank Mixing problem.

Populatrⁿ models:


$$y' = \pi y$$

$$y(0) = y_0$$



$$\left. \begin{matrix} y' = \pi y \\ y(0) = y_0 \end{matrix} \right\} \rightarrow \text{solution } y(t) = y_0 e^{\pi t}$$

exponential popⁿ growth



Logistic Growth Model

'''

$$= \pi y (1 - y/k)$$

$$y \equiv y(t)$$

Logistic Growth

$$y' = r y (1 - y/k)$$

$$y \equiv y(t)$$
$$y' \equiv y'(t)$$

Notice that there are 2 population 'states' where we have no change $y' = 0$.

$$r y (1 - y/k) = 0$$

$$y = 0$$

or

$$y = k$$

Populations where equilibrium is achieved

Goal lecture :

- ① find the closed form solution to $y' = r y (1 - y/k)$
- ② Numerical methods to solve ODE

$$\begin{aligned} y' &= r y (1 - y/k) \\ y(0) &= y_0 \end{aligned}$$

use partial fraction decomposition

step 1: $y' = \frac{dy}{dt} = r y (1 - y/k)$

$$\int \frac{dy}{y(1-y/k)} = \int r dt$$

Note: holds if $1 - y/k \neq 0$ and $y \neq 0$.

$$\int \frac{dy}{y(1-y/k)} = \int r dt$$

$$\int \frac{1}{y(1-y/k)} = \frac{A}{y} + \frac{B}{1-y/k} \Rightarrow \text{multiply both sides by } y(1-y/k)$$

$$1 = A(1-y/k) + By$$

$$y=k \Rightarrow B = 1/k$$

$$y=0 \Rightarrow A = 1$$

$$\int \frac{dy}{y(1-y/k)} = \int \left(\frac{A}{y} + \frac{B}{1-y/k} \right) dy$$

$$= \int \left(\frac{1}{y} + \frac{1/k}{1-y/k} \right) dy$$

$$= \int \frac{1}{y} dy + \frac{1}{k} \int \frac{dy}{1-y/k}$$

$$= \ln|y| - \int \frac{du}{u} \quad \begin{array}{l} \hookrightarrow \text{let } u = 1-y/k \\ du = -dy/k \end{array}$$

$$= \ln|y| - \ln|1-y/k|$$

$$= \ln \left| \frac{y}{1-y/k} \right| \quad \ln A - \ln B = \ln \left(\frac{A}{B} \right)$$

$$\int \frac{dy}{y(1-y/k)} = \pi \int dt$$

$$\ln \left| \frac{y}{1-y/k} \right| = \pi t + c$$

$$\ln \left| \frac{y}{1-y/k} \right| = e^{\pi t + c}$$

$$\ln \left| \frac{y}{1-y/k} \right| = e^{\pi t + c}$$

$$\left| \frac{y}{1-y/k} \right| = e^{\pi t + c}$$

Assume $y > 0$ $1-y/k > 0$

$$\frac{y}{1-y/k} = e^{\pi t + c}$$

$C = ?$ $y(0) = y_0$

$$\frac{y_0}{1-y_0/k} = e^{\pi \cdot 0 + c}$$

$$\frac{y_0}{1-y_0/k} = e^c$$

$$\frac{y}{1-y/k} = e^{\pi t} \cdot e^c = e^{\pi t} \cdot \frac{y_0}{1-y_0/k}$$

$$y(t) = \boxed{?}$$

currently, $\frac{y}{1-y/k} = e^{\pi t} \cdot \frac{y_0}{1-y_0/k}$

mult. by $(1-y/k)$

$$y = e^{\pi t} \frac{y_0}{1-y_0/k} (1-y/k)$$

$$y' = e^{-\pi t} \frac{y_0}{1-y_0/k} (1-y/k)$$

$$\boxed{y} = e^{\pi t} \frac{y_0}{1-y_0/k} - \frac{y e^{\pi t} y_0}{k(1-y_0/k)}$$

add $\frac{y e^{\pi t} y_0}{k(1-y_0/k)}$ to both sides.

$$\left(1 + \frac{e^{\pi t} y_0}{k(1-y_0/k)}\right) y = \frac{e^{\pi t} y_0}{1-y_0/k}$$

(k-y₀)

$$\left(1 + \frac{e^{\pi t} y_0}{k-y_0}\right) y = \frac{e^{\pi t} y_0}{1-y_0/k} = \frac{k e^{\pi t} y_0}{k-y_0}$$

$$y(t) = \frac{k e^{\pi t} y_0 / k - y_0}{1 + e^{\pi t} y_0 / k - y_0} = \frac{k y_0}{y_0 + (k - y_0) e^{-\pi t}}$$

Parameter estimation

$$y(t) = \frac{k y_0}{k + (k - y_0) e^{-\pi t}}$$

t	y(t)
1900	1
1910	10
1920	20
⋮	⋮

Numerical Solutions to ODE: Gompertz equation

$$y' = -\pi \ln\left(\frac{y}{k}\right) y$$

$$y(0) = y_0$$

$$y(0) = y_0$$

$r \rightarrow$ rate of growth $k \rightarrow$ carrying capacity of popⁿ

Numerical method to solve:

$$y' = -r \ln(y/k) y$$

Euler Method: Replace $y'(t) \rightarrow D_h y(t)$, $h = \text{timestep}$
 $= \frac{y(t+h) - y(t)}{h}$

$$\frac{y(t+h) - y(t)}{h} = -r \ln\left(\frac{y(t)}{k}\right) y(t)$$

Solve for y at time $t+h$ given $y(t)$.
 $y(t+h) = y(t) + h * \left(-r \ln\left(\frac{y(t)}{k}\right) y(t)\right)$