

Chapter Four: Integration

4.1 Antiderivatives and Indefinite Integration

Definition of Antiderivative – A function F is an antiderivative of f on an interval I if $F'(x) = f(x)$ for all x in I .

Representation of Antiderivatives – If F is an antiderivative of f on an interval I , then G is an antiderivative of f on the interval I if and only if G is of the form $G(x) = F(x) + C$, for all x in I where C is a constant.

This tells us that all antiderivatives of a function differ only by a constant C .

Examples: Find an antiderivative and then find the general antiderivative.

1. $y = 3$

A possible anti-der
is $Y = 3x$. In
general, the anti-der
is $Y = 3x + C$

2. $f(x) = 2x$

A possible $F(x)$ is
 $F(x) = x^2$. In general,
 $F(x) = x^2 + C$

3. $f(x) = 5x^4$

A possible antiderivative
is $F(x) = x^5 + 7$. In
general, $F(x) = x^5 + C$.

Since any constant has derivative 0, we use C for an arbitrary constant in the antiderivative

Notation: If we take the differential form of a derivative, $\frac{dy}{dx} = f(x)$, and rewrite it in the form

$dy = f(x)dx$ we can find the antiderivative of both sides using the integration symbol \int . That is,

$$y = \int dy = \int f(x)dx = F(x) + C$$

Each piece of this equation has a name that I will refer to: The integrand is $f(x)$, the variable of integration is given by dx , the antiderivative of $f(x)$ is $F(x)$, and the constant of integration is C . The term indefinite integral is a synonym for antiderivative.

Note: Differentiation and anti-differentiation are "inverse" operations of each other. That is, if you find the antiderivative of a function f , then take the derivative, you will end up back at f . Similarly, if you take the derivative, the antiderivative takes you back.

Some Basic Integration Rules:

$$\int 0 dx = C \qquad \int k dx = kx + C \qquad \int kf(x) dx = k \int f(x) dx$$

$$\int [f(x) \pm g(x)] dx = \int f(x) dx \pm \int g(x) dx \qquad \int x^n dx = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1 \quad \text{*important distinction!}$$

We can also consider all the trig derivatives and go backwards to find their integrals.

Examples: For each function, rewrite then integrate and finally simplify.

$$1. \int \sqrt[3]{x} dx = \int x^{1/3} dx = \frac{x^{1/3+1}}{1/3+1} + C = \frac{x^{4/3}}{4/3} + C = \frac{3}{4} x^{4/3} + C$$

$\sqrt[3]{x} = x^{1/3}$

$$2. \int \frac{1}{4x^2} dx = \int \frac{1}{4} x^{-2} dx = \frac{1}{4} \int x^{-2} dx = \frac{1}{4} \frac{x^{-2+1}}{-2+1} + C = \frac{1}{4} \frac{x^{-1}}{-1} + C = \frac{x^{-1}}{-4} + C = -\frac{1}{4x} + C$$

$$3. \int \frac{1}{x\sqrt{x}} dx = \int x^{-3/2} dx = \frac{x^{-3/2+1}}{-3/2+1} + C = \frac{x^{-1/2}}{-1/2} + C = -2x^{-1/2} + C = -\frac{2}{\sqrt{x}} + C$$

simplify $\boxed{x\sqrt{x} = x \cdot x^{1/2} = x^{1+1/2} = x^{3/2}}$

$$4. \int x(x^3+1) dx = \int (x^4+x) dx = \frac{x^{4+1}}{4+1} + \frac{x^{1+1}}{1+1} + C = \frac{x^5}{5} + \frac{x^2}{2} + C$$

No product rule for integrals!

$$5. \int \frac{1}{(3x)^2} dx = \int \frac{1}{9x^2} dx = \frac{1}{9} \int x^{-2} dx = \frac{1}{9} \frac{x^{-2+1}}{-2+1} + C = \frac{1}{9} \frac{x^{-1}}{-1} + C = -\frac{1}{9x} + C$$

on the homework:

$$6. \int \frac{1}{x^5 \sqrt{x}} dx = \int x^{-6/5} dx = \frac{x^{-6/5+1}}{-6/5+1} + C = \frac{x^{-1/5}}{-1/5} + C = -5x^{-1/5} + C = -\frac{5}{\sqrt[5]{x}} + C$$

WebAssign makes it look

weird, but this is $x \cdot \sqrt[5]{x}$

which is $x \cdot x^{1/5} = x^{6/5}$

Examples: Find the indefinite integral and check the result by differentiation.

$$1. \int (12-x) dx = 12x - \frac{x^2}{2} + C$$

$$\text{check: } \frac{d}{dx} \left(12x - \frac{x^2}{2} + C \right) = 12 - \frac{2x}{2} + 0 = 12 - x \quad \checkmark$$

$$2. \int (8x^3 - 9x^2 + 4) dx = \frac{8x^4}{4} - \frac{9x^3}{3} + 4x + C = 2x^4 - 3x^3 + 4x + C$$

$$\text{check: } \frac{d}{dx} (2x^4 - 3x^3 + 4x + C) = 8x^3 - 9x^2 + 4 + 0 = 8x^3 - 9x^2 + 4 \quad \checkmark$$

$$3. \int \left(\sqrt{x} + \frac{1}{2\sqrt{x}} \right) dx = \int \left(x^{1/2} + \frac{1}{2} x^{-1/2} \right) dx = \frac{x^{1/2+1}}{1/2+1} + \frac{1}{2} \frac{x^{-1/2+1}}{-1/2+1} + C$$

Rewrite

$$\sqrt{x} + \frac{1}{2\sqrt{x}} = x^{1/2} + \frac{1}{2} x^{-1/2}$$

$$= \frac{x^{3/2}}{3/2} + \frac{1}{2} \frac{x^{1/2}}{1/2} + C = \frac{2}{3} x^{3/2} + x^{1/2} + C$$

$$\text{check: } \frac{d}{dx} \left(\frac{2}{3} x^{3/2} + x^{1/2} + C \right) = \frac{2}{3} \left(\frac{3}{2} x^{1/2} \right) + \frac{1}{2} x^{-1/2} + 0 = \sqrt{x} + \frac{1}{2\sqrt{x}} \quad \checkmark$$

$$4. \int \frac{x^2 + 2x - 3}{x^4} dx = \int (x^{-2} + 2x^{-3} - 3x^{-4}) dx = \frac{x^{-1}}{-1} + 2 \frac{x^{-2}}{-2} - \frac{3x^{-3}}{-3} + C = -x^{-1} - x^{-2} + x^{-3} + C$$

$$= -\frac{1}{x} - \frac{1}{x^2} + \frac{1}{x^3} + C$$

NO quotient rule for integrals!

$$\frac{x^2 + 2x - 3}{x^4} = \frac{x^2}{x^4} + \frac{2x}{x^4} - \frac{3}{x^4} = x^{-2} + 2x^{-3} - 3x^{-4}$$

$$\text{Check: } \frac{d}{dx} (-x^{-1} - x^{-2} + x^{-3} + C) = -(-1x^{-2}) - (-2x^{-3}) + (-3x^{-4}) + 0 = x^{-2} + 2x^{-3} - 3x^{-4} \quad \checkmark$$

$$5. \int (2t^2 - 1)^2 dt = \int (4t^4 - 4t^2 + 1) dt = 4 \frac{t^5}{5} - 4 \frac{t^3}{3} + t + C = \frac{4}{5} t^5 - \frac{4}{3} t^3 + t + C$$

Rewrite $(2t^2 - 1)^2 = 4t^4 - 4t^2 + 1$

$$\text{Check: } \frac{d}{dt} \left(\frac{4}{5} t^5 - \frac{4}{3} t^3 + t + C \right) = \frac{4}{5} (5t^4) - \frac{4}{3} (3t^2) + 1 + 0 = 4t^4 - 4t^2 + 1 \quad \checkmark$$

$$6. \int (t^2 - \cos t) dt = \int t^2 dt - \int \cos t dt = \frac{t^3}{3} - \sin t + C$$

$$\text{Check: } \frac{d}{dt} \left(\frac{t^3}{3} - \sin t + C \right) = \frac{1}{3} (3t^2) - \cos t + 0 = t^2 - \cos t \quad \checkmark$$

$$7. \int (\theta^2 + \sec^2 \theta) d\theta = \frac{\theta^3}{3} + \tan \theta + C$$

↑
What has derivative equal to $\sec^2 \theta$?

$$\text{Check: } \frac{d}{d\theta} \left(\frac{\theta^3}{3} + \tan \theta + C \right) = \frac{3\theta^2}{3} + \sec^2 \theta + 0 = \theta^2 + \sec^2 \theta \quad \checkmark$$

$$8. \int \sec y (\tan y - \sec y) dy = \int (\sec y \tan y - \sec^2 y) dy = \sec y - \tan y + C$$

still no product rule!

$$\text{Check: } \frac{d}{dy} (\sec y - \tan y + C) = \sec y \tan y - \sec^2 y + 0 = \sec y \tan y - \sec^2 y \quad \checkmark$$

Example: Find the equation of y given $\frac{dy}{dx} = 2x - 1$ that has the particular point $(1, 1)$ as part of its solution set.

$$y = \int dy = \int (2x - 1) dx = 2 \frac{x^2}{2} - x + C = x^2 - x + C$$

Since $(1, 1)$ is a solution, we can substitute to find a specific value of C :

$$1 = (1)^2 - (1) + C$$

$$1 = 1 - 1 + C$$

$$1 = C$$

$$\text{Solution: } y = x^2 - x + 1$$

Example: Solve the differential equation.

1. $f'(x) = 6x^2, f(0) = -1$

$$f(x) = \int 6x^2 dx = 6 \frac{x^3}{3} + C = 2x^3 + C$$

$$-1 = 2(0)^3 + C$$

$$-1 = C$$

$$f(x) = 2x^3 - 1$$

2. $f'(p) = 10p - 12p^3, f(3) = 2$

$$f(p) = \int (10p - 12p^3) dp = \frac{10p^2}{2} - \frac{12p^4}{4} + C = 5p^2 - 3p^4 + C$$

$$2 = 5(3)^2 - 3(3)^4 + C$$

$$2 = 45 - 243 + C$$

$$200 = C$$

$$f(p) = 5p^2 - 3p^4 + 200$$

3. $h''(x) = \sin x, h'(0) = 1, h(0) = 6$

$$h'(x) = \int h''(x) dx = \int \sin x dx = -\cos x + C$$

$$1 = -\cos(0) + C$$

$$1 = -1 + C$$

$$2 = C$$

$$h'(x) = -\cos x + 2$$

$$h(x) = \int (-\cos x + 2) dx = -\sin x + 2x + C$$

$$6 = -\sin(0) + 2(0) + C$$

$$6 = C$$

$$h(x) = -\sin x + 2x + 6$$

Example: A particle, initially at rest, moves along the x -axis such that its acceleration at time $t > 0$ is given by $a(t) = \cos t$. At the time $t = 0$, its position is $x = 3$.

Similar to
#8 on homework

- a) Find the velocity and position functions for the particle.
b) Find the values of t for which the particle is at rest.

$$a(t) = \cos t$$

$$v(t) = \int a(t) dt = \int \cos t dt = \sin t + C$$

at rest has $v(0) = 0$
so $C = 0$

part a → $v(t) = \sin t$

$$x(t) = \int v(t) dt = \int \sin t dt = -\cos t + C$$

$$\begin{aligned} 3 &= -\cos(0) + C \\ 3 &= -1 + C \\ 4 &= C \end{aligned}$$

part a → $x(t) = -\cos t + 4$

b) particle at rest \Rightarrow velocity is 0

$$v(t) = \sin t = 0 \quad \text{when } t = k\pi \quad \text{for any integer } k$$