

# A non-partitionable Cohen-Macaulay simplicial complex

Art Duval<sup>1</sup>, Bennet Goeckner<sup>2</sup>, Caroline Klivans<sup>3</sup>,  
Jeremy Martin<sup>2</sup>

<sup>1</sup>University of Texas at El Paso, <sup>2</sup>University of Kansas, <sup>3</sup>Brown University

Discrete Geometry and Algebraic Combinatorics Conference  
University of Texas Rio Grande Valley  
May 4, 2016

Preprint: <http://arxiv.org/abs/1504.04279>

# Partitionability Conjecture

Richard Stanley: "... a central combinatorial conjecture on Cohen-Macaulay complexes is the following."

Conjecture (Stanley '79; Garsia '80)

*Every Cohen-Macaulay simplicial complex is partitionable.*

# Partitionability Conjecture

Richard Stanley: "... a central combinatorial conjecture on Cohen-Macaulay complexes is the following."

Conjecture (Stanley '79; Garsia '80)

*Every Cohen-Macaulay simplicial complex is partitionable.*

Theorem (DGKM '15)

*No.*

# Partitionability Conjecture

Richard Stanley: "... a central combinatorial conjecture on Cohen-Macaulay complexes is the following."

Conjecture (Stanley '79; Garsia '80)

*Every Cohen-Macaulay simplicial complex is partitionable.*

Theorem (DGKM '15)

*No.*

Stanley: "I am glad that this problem has finally been put to rest,

# Partitionability Conjecture

Richard Stanley: "... a central combinatorial conjecture on Cohen-Macaulay complexes is the following."

Conjecture (Stanley '79; Garsia '80)

*Every Cohen-Macaulay simplicial complex is partitionable.*

Theorem (DGKM '15)

*No.*

Stanley: "I am glad that this problem has finally been put to rest, though I would have preferred a proof rather than a counterexample."

# Partitionability Conjecture

Richard Stanley: "... a central combinatorial conjecture on Cohen-Macaulay complexes is the following."

Conjecture (Stanley '79; Garsia '80)

*Every Cohen-Macaulay simplicial complex is partitionable.*

Theorem (DGKM '15)

*No.*

Stanley: "I am glad that this problem has finally been put to rest, though I would have preferred a proof rather than a counterexample. Perhaps you can withdraw your paper from the arXiv and come up with a proof instead."

# Counting faces of spheres

## Definition (Sphere)

Simplicial complex whose realization is a triangulation of a sphere.

# Counting faces of spheres

## Definition (Sphere)

Simplicial complex whose realization is a triangulation of a sphere.

## Definition ( $f$ -vector)

$f_i = f_i(\Delta) =$  number of  $i$ -dimensional faces of  $\Delta$ .

## Conjecture (Upper Bound)

*Explicit upper bound on  $f_i$  of a sphere with given dimension and number of vertices.*



# Counting faces of spheres

## Definition (Sphere)

Simplicial complex whose realization is a triangulation of a sphere.

## Definition ( $f$ -vector)

$f_i = f_i(\Delta) =$  number of  $i$ -dimensional faces of  $\Delta$ .

## Conjecture (Upper Bound)

*Explicit upper bound on  $f_i$  of a sphere with given dimension and number of vertices.*

This was proved by Stanley in 1975. Some of the key ingredients:

- ▶ face-ring (algebraic object derived from the simplicial complex) [Stanley, Hochster]
- ▶ face-ring of sphere is **Cohen-Macaulay** [Reisner]

# Cohen-Macaulay simplicial complexes

CM *rings* of great interest in commutative algebra (depth = dimension). Here is a more topological/combinatorial definition.

## Definition (Link)

$\text{lk}_\Delta \sigma = \{\tau \in \Delta : \tau \cap \sigma = \emptyset, \tau \cup \sigma \in \Delta\}$ , what  $\Delta$  looks like near  $\sigma$ .

# Cohen-Macaulay simplicial complexes

CM *rings* of great interest in commutative algebra (depth = dimension). Here is a more topological/combinatorial definition.

## Definition (Link)

$\text{lk}_\Delta \sigma = \{\tau \in \Delta : \tau \cap \sigma = \emptyset, \tau \cup \sigma \in \Delta\}$ , what  $\Delta$  looks like near  $\sigma$ .

## Definition (Homology)

$\tilde{H}_i(\Delta) = \ker \partial_i / \text{im } \partial_{i+1}$ , measures  $i$ -dimensional “holes” of  $\Delta$ .

# Cohen-Macaulay simplicial complexes

CM *rings* of great interest in commutative algebra (depth = dimension). Here is a more topological/combinatorial definition.

## Definition (Link)

$\text{lk}_\Delta \sigma = \{\tau \in \Delta : \tau \cap \sigma = \emptyset, \tau \cup \sigma \in \Delta\}$ , what  $\Delta$  looks like near  $\sigma$ .

## Definition (Homology)

$\tilde{H}_i(\Delta) = \ker \partial_i / \text{im } \partial_{i+1}$ , measures  $i$ -dimensional “holes” of  $\Delta$ .

## Theorem (Reisner '76)

Face-ring of  $\Delta$  is *Cohen-Macaulay* if, for all  $\sigma \in \Delta$ ,

$$\tilde{H}_i(\text{lk}_\Delta \sigma) = 0 \quad \text{for } i < \dim \text{lk}_\Delta \sigma.$$

We take this as our definition of CM simplicial complex.

# Cohen-Macaulayness is topological

Recall our definition:

Theorem (Reisner '76)

Face-ring of  $\Delta$  is *Cohen-Macaulay* if, for all  $\sigma \in \Delta$ ,

$$\tilde{H}_i(\text{lk}_\Delta \sigma) = 0 \quad \text{for } i < \dim \text{lk}_\Delta \sigma.$$

# Cohen-Macaulayness is topological

Recall our definition:

Theorem (Reisner '76)

Face-ring of  $\Delta$  is *Cohen-Macaulay* if, for all  $\sigma \in \Delta$ ,

$$\tilde{H}_i(\text{lk}_\Delta \sigma) = 0 \quad \text{for } i < \dim \text{lk}_\Delta \sigma.$$

Munkres ('84) showed that CM is a **topological** condition. That is, it only depends on (the homeomorphism class of) the **realization** of  $\Delta$ . In particular, spheres and balls are CM.

Example



is **not** CM

The conditions for the UBC most easily stated in terms of *h*-vector.

## Definition (*h*-vector)

Let  $\dim \Delta = d - 1$ .

$$\sum_{i=0}^d f_{i-1}(t-1)^{d-i} = \sum_{k=0}^d h_k t^{d-k}$$

The *h*-vector of  $\Delta$  is  $h(\Delta) = (h_0, h_1, \dots, h_d)$ . Coefficients not always non-negative!

The conditions for the UBC most easily stated in terms of *h*-vector.

## Definition (*h*-vector)

Let  $\dim \Delta = d - 1$ .

$$\sum_{i=0}^d f_{i-1}(t-1)^{d-i} = \sum_{k=0}^d h_k t^{d-k}$$


Equivalently,

$$\sum_{i=0}^d f_{i-1} t^{d-i} = \sum_{k=0}^d h_k (t+1)^{d-k}$$

The *h*-vector of  $\Delta$  is  $h(\Delta) = (h_0, h_1, \dots, h_d)$ . Coefficients not always non-negative!



# Example




$f(\Delta) = (1, 5, 9, 6)$ , and

$$1t^3 + 5t^2 + 9t + 6 = 1(t+1)^3 + 2(t+1)^2 + 2(t+1)^1 + 1$$

so  $h(\Delta) = (1, 2, 2, 1)$ .

# Example



$f(\Delta) = (1, 5, 9, 6)$ , and


$$1t^3 + 5t^2 + 9t + 6 = 1(t+1)^3 + 2(t+1)^2 + 2(t+1)^1 + 1$$

so  $h(\Delta) = (1, 2, 2, 1)$ .

Note that in this case,  $h \geq 0$ . This is a consequence of the algebraic defn of CM. But how could we see this combinatorially?


# Partitionability

$$1t^3 + 5t^2 + 9t + 6 = 1(t+1)^3 + 2(t+1)^2 + 2(t+1)^1 + 1$$



# Partitionability

$$1t^3 + 5t^2 + 9t + 6 = 1(t+1)^3 + 2(t+1)^2 + 2(t+1)^1 + 1$$



## Definition (Partitionable)

When a simplicial complex can be **partitioned** like this, into Boolean intervals whose tops are facets, we say the complex is **partitionable**.

# Shellability

Most CM complexes in combinatorics are shellable:

## Definition (Shellable)

A simplicial complex is **shellable** if it can be built one facet at a time, so that there is always a unique new minimal face being added.

A shelling is a particular kind of partitioning.

# Shellability

Most CM complexes in combinatorics are shellable:

## Definition (Shellable)

A simplicial complex is **shellable** if it can be built one facet at a time, so that there is always a unique new minimal face being added.

A shelling is a particular kind of partitioning.

## Proposition

*If  $\Delta$  is shellable, then  $h_k$  counts number of intervals whose bottom (the unique new minimal face) is dimension  $k - 1$ .*

## Example

In our previous example, minimal new faces were:  $\emptyset$ , **vertex**, **edge**, **vertex**, **edge**, triangle.

# Relative simplicial complexes


## Definition (Relative simplicial complex)

$\Phi$  is a **relative simplicial complex** on  $V$  if:

- ▶  $\Phi \subseteq 2^V$ ; and
- ▶  $\rho \subseteq \sigma \subseteq \tau$  and  $\rho, \tau \in \Phi$  together imply  $\sigma \in \Phi$

We can write any relative complex  $\Phi$  as  $\Phi = (\Delta, \Gamma)$ , for some pair of simplicial complexes  $\Gamma \subseteq \Delta$ .

## Example



# Relative simplicial complexes


## Definition (Relative simplicial complex)

$\Phi$  is a **relative simplicial complex** on  $V$  if:

- ▶  $\Phi \subseteq 2^V$ ; and
- ▶  $\rho \subseteq \sigma \subseteq \tau$  and  $\rho, \tau \in \Phi$  together imply  $\sigma \in \Phi$

We can write any relative complex  $\Phi$  as  $\Phi = (\Delta, \Gamma)$ , for some pair of simplicial complexes  $\Gamma \subseteq \Delta$ . But  $\Delta$  and  $\Gamma$  are not unique.

## Example





# Relative Cohen-Macaulay

Recall  $\Delta$  is CM when

$$\tilde{H}_i(\mathrm{lk}_\Delta \sigma) = 0 \quad \text{for } i < \mathrm{lk}_\Delta \sigma.$$

# Relative Cohen-Macaulay

Recall  $\Delta$  is CM when

$$\tilde{H}_i(\mathrm{lk}_\Delta \sigma) = 0 \quad \text{for } i < \mathrm{lk}_\Delta \sigma.$$

This generalizes easily:

**Theorem (Stanley '87)**

Face-ring of  $\Phi = (\Delta, \Gamma)$  is *relative Cohen-Macaulay* if, for all  $\sigma \in \Delta$ ,

$$\tilde{H}_i(\mathrm{lk}_\Delta \sigma, \mathrm{lk}_\Gamma \sigma) = 0 \quad \text{for } i < \mathrm{lk}_\Delta \sigma.$$

# Relative Cohen-Macaulay

Recall  $\Delta$  is CM when

$$\tilde{H}_i(\mathrm{lk}_\Delta \sigma) = 0 \quad \text{for } i < \mathrm{lk}_\Delta \sigma.$$


This generalizes easily:

**Theorem (Stanley '87)**

Face-ring of  $\Phi = (\Delta, \Gamma)$  is *relative Cohen-Macaulay* if, for all  $\sigma \in \Delta$ ,

$$\tilde{H}_i(\mathrm{lk}_\Delta \sigma, \mathrm{lk}_\Gamma \sigma) = 0 \quad \text{for } i < \mathrm{lk}_\Delta \sigma.$$

**Example**



# Looking for a non-trivial example

We were trying to **prove** the conjecture.

# Looking for a non-trivial example

We were trying to **prove** the conjecture. Idea of our “proof”:

- ▶ Remove all the faces containing a given vertex (this will be the first part of the partitioning).

# Looking for a non-trivial example

We were trying to **prove** the conjecture. Idea of our “proof”:

- ▶ Remove all the faces containing a given vertex (this will be the first part of the partitioning).
- ▶ Try to make sure what's left is **relative** CM.

# Looking for a non-trivial example

We were trying to **prove** the conjecture. Idea of our “proof”:

- ▶ Remove all the faces containing a given vertex (this will be the first part of the partitioning).
- ▶ Try to make sure what's left is **relative** CM.
- ▶ Apply induction.

# Looking for a non-trivial example

We were trying to **prove** the conjecture. Idea of our “proof”:

- ▶ Remove all the faces containing a given vertex (this will be the first part of the partitioning).
- ▶ Try to make sure what's left is **relative** CM.
- ▶ Apply induction.
- ▶ How hard is it to take that **second** step of the partitioning, which is the first step for the relative complex?



# Looking for a non-trivial example

We were trying to **prove** the conjecture. Idea of our “proof”:

- ▶ Remove all the faces containing a given vertex (this will be the first part of the partitioning).
- ▶ Try to make sure what's left is **relative** CM.
- ▶ Apply induction.
- ▶ How hard is it to take that **second** step of the partitioning, which is the first step for the relative complex?
- ▶ We wanted to find a non-trivial example of something CM and partitionable, so we could see how it would work.



# Looking for a non-trivial example

We were trying to **prove** the conjecture. Idea of our “proof”:

- ▶ Remove all the faces containing a given vertex (this will be the first part of the partitioning).
- ▶ Try to make sure what's left is **relative** CM.
- ▶ Apply induction.
- ▶ How hard is it to take that **second** step of the partitioning, which is the first step for the relative complex?
- ▶ We wanted to find a non-trivial example of something CM and partitionable, so we could see how it would work.
- ▶ Idea: non-trivial = not shellable; CM = ball (and if it's not partitionable, we're done). So we are looking for **non-shellable balls**.



# Ziegler's non-shellable ball ('98)

Non-shellable 3-ball with 10 vertices and 21 tetrahedra



# Ziegler's non-shellable ball ('98)

Non-shellable 3-ball with 10 vertices and 21 tetrahedra




Just because it is partitionable does not mean you can start partitioning in any order.

So we started to partition until we could not go any further (without backtracking). This part uses the computer!

# First pass with Ziegler

We found a relative complex  $Q_5 = (X_5, A_5)$


- ▶  $X_5$  has 6 vertices, 5 facets



# First pass with Ziegler

We found a relative complex  $Q_5 = (X_5, A_5)$


- ▶  $X_5$  has 6 vertices, 5 facets
- ▶ remove  $A_5$ , which is 4 triangles on boundary



# First pass with Ziegler

We found a relative complex  $Q_5 = (X_5, A_5)$


- ▶  $X_5$  has 6 vertices, 5 facets
- ▶ remove  $A_5$ , which is 4 triangles on boundary
- ▶ relative CM (since  $X_5$  and  $A_5$  shellable,  $A_5 \subseteq \partial X_5$ )



# First pass with Ziegler

We found a relative complex  $Q_5 = (X_5, A_5)$

- ▶  $X_5$  has 6 vertices, 5 facets
- ▶ remove  $A_5$ , which is 4 triangles on boundary
- ▶ relative CM (since  $X_5$  and  $A_5$  shellable,  $A_5 \subseteq \partial X_5$ )
- ▶ not partitionable






# First pass with Ziegler

We found a relative complex  $Q_5 = (X_5, A_5)$


- ▶  $X_5$  has 6 vertices, 5 facets
- ▶ remove  $A_5$ , which is 4 triangles on boundary
- ▶ relative CM (since  $X_5$  and  $A_5$  shellable,  $A_5 \subseteq \partial X_5$ )
- ▶ not partitionable



# First pass with Ziegler

We found a relative complex  $Q_5 = (X_5, A_5)$


- ▶  $X_5$  has 6 vertices, 5 facets
- ▶ remove  $A_5$ , which is 4 triangles on boundary
- ▶ relative CM (since  $X_5$  and  $A_5$  shellable,  $A_5 \subseteq \partial X_5$ )
- ▶ not partitionable



# First pass with Ziegler

We found a relative complex  $Q_5 = (X_5, A_5)$

- ▶  $X_5$  has 6 vertices, 5 facets
- ▶ remove  $A_5$ , which is 4 triangles on boundary
- ▶ relative CM (since  $X_5$  and  $A_5$  shellable,  $A_5 \subseteq \partial X_5$ )
- ▶ not partitionable



If only we could build a non-relative complex out of this.

## Proposition

*If  $X$  and  $(X, A)$  are CM and  $\dim A = \dim X - 1$ , then gluing together two copies of  $X$  along  $A$  gives a CM (non-relative) complex.*

If we glue together **two** copies of  $X$  along  $A$ , is it partitionable?

## Proposition

*If  $X$  and  $(X, A)$  are CM and  $\dim A = \dim X - 1$ , then gluing together two copies of  $X$  along  $A$  gives a CM (non-relative) complex.*

If we glue together **two** copies of  $X$  along  $A$ , is it partitionable?  
Maybe: some parts of  $A$  can help partition one copy of  $X$ , other parts of  $A$  can help partition the other copy of  $X$ .

# Pigeonhole principle

Recall our example  $(X, A)$  is:

- ▶ relative Cohen-Macaulay
- ▶ not partitionable

## Remark

If we glue together **many** copies of  $X$  along  $A$ , at least one copy will be missing all of  $A$ !

# Pigeonhole principle

Recall our example  $(X, A)$  is:

- ▶ relative Cohen-Macaulay
- ▶ not partitionable

## Remark

If we glue together **many** copies of  $X$  along  $A$ , at least one copy will be missing all of  $A$ ! How many is enough?

# Pigeonhole principle

Recall our example  $(X, A)$  is:

- ▶ relative Cohen-Macaulay
- ▶ not partitionable

## Remark

If we glue together **many** copies of  $X$  along  $A$ , at least one copy will be missing all of  $A$ ! How many is enough? More than the number of all faces in  $A$ . Then the result will **not** be partitionable.



# Pigeonhole principle

Recall our example  $(X, A)$  is:

- ▶ relative Cohen-Macaulay
- ▶ not partitionable

## Remark

If we glue together **many** copies of  $X$  along  $A$ , at least one copy will be missing all of  $A$ ! How many is enough? More than the number of all faces in  $A$ . Then the result will **not** be partitionable.

## Remark

But the resulting complex is not actually a simplicial complex because of repeats.

# Pigeonhole principle

Need our example  $(X, A)$  to be:

- ▶ relative Cohen-Macaulay
- ▶ not partitionable
- ▶  $A$  **vertex-induced** (minimal faces of  $(X, A)$  are vertices)

## Remark

If we glue together **many** copies of  $X$  along  $A$ , at least one copy will be missing all of  $A$ ! How many is enough? More than the number of all faces in  $A$ . Then the result will **not** be partitionable.

## Remark

But the resulting complex is not actually a simplicial complex because of repeats. To avoid this problem, we need to make sure that  $A$  is **vertex-induced**. This means every face in  $X$  among vertices in  $A$  must be in  $A$  as well. (Minimal faces of  $(X, A)$  are vertices.)

# Eureka!

By computer search, we found that if

- ▶  $Z$  is Ziegler's 3-ball, and
- ▶  $B = Z$  restricted to all vertices except 1,5,9 ( $B$  has 7 facets),

then  $Q = (Z, B)$  satisfies all our criteria!


# Eureka!

By computer search, we found that if

- ▶  $Z$  is Ziegler's 3-ball, and
- ▶  $B = Z$  restricted to all vertices except 1,5,9 ( $B$  has 7 facets),

then  $Q = (Z, B)$  satisfies all our criteria!

Also  $Q = (X, A)$ , where  $X$  has 14 facets, and  $A$  is 5 triangles:



1249	1269
1569	1589
1489	1458
1457	4578
1256	0125
0256	0123
1234	1347

# Putting it all together

- ▶ Since  $A$  has 24 faces total (including the empty face), we know gluing together 25 copies of  $X$  along their common copy of  $A$ , the resulting (non-relative) complex  $C_{25}$  is CM, not partitionable.

## Putting it all together

- ▶ Since  $A$  has 24 faces total (including the empty face), we know gluing together 25 copies of  $X$  along their common copy of  $A$ , the resulting (non-relative) complex  $C_{25}$  is CM, not partitionable.
- ▶ In fact, computer search showed that gluing together only 3 copies of  $X$  will do it. Resulting complex  $C_3$  has  $f$ -vector  $(1, 16, 71, 98, 42)$ .

## Putting it all together

- ▶ Since  $A$  has 24 faces total (including the empty face), we know gluing together 25 copies of  $X$  along their common copy of  $A$ , the resulting (non-relative) complex  $C_{25}$  is CM, not partitionable.
- ▶ In fact, computer search showed that gluing together only 3 copies of  $X$  will do it. Resulting complex  $C_3$  has  $f$ -vector  $(1, 16, 71, 98, 42)$ .
- ▶ Later we found short proof by hand to show that  $C_3$  works.

# Stanley depth (a brief summary)

## Definition (Stanley)

If  $I$  is a monomial ideal in a polynomial ring  $S$ , then the **Stanley depth**  $\text{sdepth } S/I$  is a purely combinatorial analogue of depth, defined in terms of certain vector space decompositions of  $S/I$ .

## Conjecture (Stanley '82)

*For all monomial ideals  $I$ ,  $\text{sdepth } S/I \geq \text{depth } S/I$ .*

## Theorem (Herzog, Jahan, Yassemi '08)

*If  $I$  is the **Stanley-Reisner ideal** (related to the face ring) of a Cohen-Macaulay complex  $\Delta$ , then the inequality  $\text{sdepth } S/I \geq \text{depth } S/I$  is equivalent to the partitionability of  $\Delta$ .*

## Corollary

*Our counterexample disproves this conjecture as well.*



# Constructibility

## Definition

A  $d$ -dimensional simplicial complex  $\Delta$  is **constructible** if:

- ▶ it is a simplex; or
- ▶  $\Delta = \Delta_1 \cup \Delta_2$ , where  $\Delta_1, \Delta_2, \Delta_1 \cap \Delta_2$  are constructible of dimensions  $d, d, d - 1$ , respectively.

# Constructibility

## Definition

A  $d$ -dimensional simplicial complex  $\Delta$  is **constructible** if:

- ▶ it is a simplex; or
- ▶  $\Delta = \Delta_1 \cup \Delta_2$ , where  $\Delta_1, \Delta_2, \Delta_1 \cap \Delta_2$  are constructible of dimensions  $d, d, d - 1$ , respectively.

## Theorem

*Constructible complexes are Cohen-Macaulay.*

# Constructibility

## Definition

A  $d$ -dimensional simplicial complex  $\Delta$  is **constructible** if:

- ▶ it is a simplex; or
- ▶  $\Delta = \Delta_1 \cup \Delta_2$ , where  $\Delta_1, \Delta_2, \Delta_1 \cap \Delta_2$  are constructible of dimensions  $d, d, d - 1$ , respectively.

## Theorem

*Constructible complexes are Cohen-Macaulay.*

## Question (Hachimori '00)

*Are constructible complexes partitionable?*

# Constructibility

## Definition

A  $d$ -dimensional simplicial complex  $\Delta$  is **constructible** if:

- ▶ it is a simplex; or
- ▶  $\Delta = \Delta_1 \cup \Delta_2$ , where  $\Delta_1, \Delta_2, \Delta_1 \cap \Delta_2$  are constructible of dimensions  $d, d, d - 1$ , respectively.

## Theorem

*Constructible complexes are Cohen-Macaulay.*

## Question (Hachimori '00)

*Are constructible complexes partitionable?*

## Corollary

*Our counterexample is constructible, so the answer to this question is no.*

# Open question: Smaller counterexample?

Open questions:

## Question

*Is there a smaller 3-dimensional counterexample to the partitionability conjecture?*

# Open question: Smaller counterexample?

Open questions:

## Question

*Is there a smaller 3-dimensional counterexample to the partitionability conjecture?*

## Question

*Is the partitionability conjecture true in 2 dimensions?*

# Save the conjecture: Strengthen the hypothesis

More open questions (based on what our counterexample is **not**):  
Note that our counterexample is not a ball (3 balls sharing common 2-dimensional faces), but all balls are CM.

## Question

*Are simplicial **balls** partitionable?*

# Save the conjecture: Strengthen the hypothesis

More open questions (based on what our counterexample is **not**):  
Note that our counterexample is not a ball (3 balls sharing common 2-dimensional faces), but all balls are CM.

## Question

*Are simplicial **balls** partitionable?*

## Definition (Balanced)

A simplicial complex is **balanced** if vertices can be colored so that every facet has one vertex of each color.

## Question

*Are **balanced** Cohen-Macaulay complexes partitionable?*



# Save the conjecture: Weaken the conclusion

## Question

What *does* the  $h$ -vector of a CM complex count?

# Save the conjecture: Weaken the conclusion

## Question

What *does* the  $h$ -vector of a CM complex count?

One possible answer (D.-Zhang '01) replaces Boolean intervals with “Boolean trees”. But maybe there are other answers.

