

## Math 2313, Final

Name \_\_\_\_\_

1. Consider the surface  $x^3 \ln(y) + ze^{xz} = 0$ :

a. Write the equation for the tangent plane at the point  $(1, 1, 0)$ .

answer:  $y + z = 1$

b. Write the parametric equations for the normal line at  $(1, 1, 0)$ .

answer:  $(x, y, z) = (1, 1 + t, t)$

2. Consider a particle whose position is given by the vector  $r(t) = \langle \cos(3t), \sin(3t), 4t \rangle$ . Find the velocity  $r'(t)$ , speed  $\|r'(t)\|$ , and acceleration  $r''(t)$ . Show that the acceleration is always perpendicular to the velocity.

answer:  $r' = \langle -3\sin(3t), 3\cos(3t), 4 \rangle$ ,  $\|r'\| = 5$   
 $r'' = \langle -9\cos(3t), -9\sin(3t), 0 \rangle$   
 $r' \bullet r'' = 0$

3. Express  $(\frac{\partial u}{\partial r})^2 + \frac{1}{r^2}(\frac{\partial u}{\partial \theta})^2$  in terms of  $\frac{\partial u}{\partial x}$  and  $\frac{\partial u}{\partial y}$ , using the chain rule, and simplify as much as possible. (Hint:  $x = r \cos(\theta), y = r \sin(\theta)$ )

answer:  $(\frac{\partial u}{\partial x})^2 + (\frac{\partial u}{\partial y})^2$

4. Find the directional derivative of  $f(x, y, z) = x^3 + \ln(xy) + e^{3yz}$  at the point  $(1, 1, 0)$  in the direction of the vector  $\langle 1, 1, -1 \rangle$ .

answer:  $\frac{2}{\sqrt{3}}$

5. Find the dimensions of an open box (no top) with a volume of  $32\text{cm}^3$  which uses the least amount of cardboard.

answer:  $x = 4\text{cm}, y = 4\text{cm}, z = 2\text{cm}$

6. Evaluate  $\int_0^1 \int_y^1 \sqrt{1-x^2} dx dy$ . (Hint: you will need to reverse the order of integration.)

answer:  $\frac{1}{3}$

7. Write an integral which, if evaluated (but don't evaluate), would give the mass of the tetrahedron in the first octant under the plane  $4x + 2y + z = 12$ , if the density is  $\rho(x, y, z)$ .

answer:  $\int_0^3 \int_0^{6-2x} \int_0^{12-4x-2y} \rho(x, y, z) dz dy dx$